

B70

7a

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 33/50, C12Q 1/68, G01N 33/72, 33/574		A1	(11) International Publication Number: WO 99/53316
			(43) International Publication Date: 21 October 1999 (21.10.99)
(21) International Application Number: PCT/US99/08013		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 13 April 1999 (13.04.99)			
(30) Priority Data: 09/059,713 13 April 1998 (13.04.98) US		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71) Applicant: EXACT LABORATORIES, INC. [US/US]; 63 Great Road, Maynard, MA 01754 (US).			
(72) Inventors: LAPIDUS, Stanley, N.; 12 Old Evergreen Road, Bedford, NH 03110 (US). SHUBER, Anthony, P.; 11 Grant Street, Milford, MA 01757 (US).			
(74) Agent: WALLER, Patrick, R., H.; Testa, Hurwitz & Thibault, LLP, High Street Tower, 125 High Street, Boston, MA 02110 (US).			
(54) Title: METHODS FOR DISEASE DIAGNOSIS FROM STOOL SAMPLES			
(57) Abstract The present invention provides methods for preparing a stool sample in order to screen for the presence of indicators of a disease, for example a subpopulation of cancerous or precancerous cells. The methods take advantage of the recognition that cellular debris from cancerous and precancerous cells is deposited onto only a longitudinal stripe of stool as the stool is forming in the colon. Accordingly, methods of the invention comprise obtaining a representative sample, such as a circumferential or cross-sectional sample of stool in order to ensure that any disease indicator, such as cellular debris that is shed by colonic cells, is obtained in the sample.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHODS FOR DISEASE DIAGNOSIS FROM STOOL SAMPLES

This patent application is a continuation-in-part of U.S. Serial Number 08/699,678, filed August 14, 1996, the disclosure of which is incorporated herein by reference.

FIELD OF THE INVENTION

5 This invention relates to methods for disease diagnosis, including the early detection of colon cancer in patients, and more particularly to methods for preparing stool samples for disease diagnosis, including the detection of colon cancer, so as to assure or increase the likelihood that the sample will contain the diagnostically relevant information if the patient has a disease, for example a cancerous or precancerous
10 lesion, and to methods for stool sample analysis.

BACKGROUND OF THE INVENTION

 Stool samples frequently must be prepared for medical diagnostic analysis. Stool samples may be analyzed to help diagnose medical conditions ranging from parasitic, bacterial or viral infections to inflammatory bowel disease and colorectal
15 cancer.

 Colorectal cancer is a leading cause of death in Western society. However, if diagnosed early, it may be treated effectively by surgical removal of the cancerous tissue. Colorectal cancers originate in the colorectal epithelium and typically are not extensively vascularized (and therefore not invasive) during the early stages of
20 development. Colorectal cancer is thought to result from the clonal expansion of a single mutant cell in the epithelial lining of the colon or rectum. The transition to a highly vascularized, invasive and ultimately metastatic cancer which spreads throughout the body commonly takes ten years or longer. If the cancer is detected prior to invasion, surgical removal of the cancerous tissue is an effective cure.
25 However, colorectal cancer is often detected only upon manifestation of clinical symptoms, such as pain and black tarry stool. Generally, such symptoms are present only when the disease is well established, often after metastasis has occurred, and the

- 2 -

prognosis for the patient is poor, even after surgical resection of the cancerous tissue. Early detection of colorectal cancer therefore is important in that detection may significantly reduce its morbidity.

Invasive diagnostic methods such as endoscopic examination allow for direct
5 visual identification, removal, and biopsy of potentially cancerous growths such as polyps. Endoscopy is expensive, uncomfortable, inherently risky, and therefore not a practical tool for screening populations to identify those with colorectal cancer. Non-invasive analysis of stool samples for characteristics indicative of the presence of colorectal cancer or precancer is a preferred alternative for early diagnosis, but no
10 known diagnostic method is available which reliably achieves this goal.

Current non-invasive diagnostic methods involve assaying stool samples for the presence of fecal occult blood or for elevated levels of carcinoembryonic antigen, both of which are suggestive of the presence of colorectal cancer. Additionally, recent developments in molecular biology provide methods of great potential for detecting the
15 presence of a range of DNA mutations or alterations associated with and indicative of the presence of colorectal cancer. The presence of such mutations theoretically can be detected in DNA found in stool samples during the early stages of colorectal cancer. However, stool comprises cells and cellular debris from the patient, from microorganisms, and from food, resulting in a heterogeneous population of cells. This
20 makes detection of a small, specific subpopulation impossible to detect reliably.

Stool diagnostic assays for colorectal cancer described in the art typically are performed on samples prepared from randomly sampled portions of voided stool. However, samples prepared according to such methods do not reproducibly yield characteristics indicative of the presence of colorectal cancer or precancer, even when
25 prepared from stool voided by a patient with colorectal cancer or precancer. There is therefore a need in the art for methods for early diagnosis of colorectal cancer or precancer that will reproducibly detect characteristics indicative of the presence of cancerous or precancerous material in samples prepared from stool voided by a patient with colorectal cancer or precancer. Such methods are provided herein.

- 3 -

SUMMARY OF THE INVENTION

It has now been appreciated that cells and cellular debris are shed from colonic epithelial cells onto forming stool in a longitudinal "stripe" of material along the length of the stool. The shed material is confined to this longitudinal stripe, as shown in Figure 1 (designated "C"). Based upon this recognition, Applicants teach that stool sample preparation for diagnostic testing must include taking a representative sample in order to ensure that the sample will contain any cells or cellular debris that was shed into the stool as it passed through the colon. Accordingly, methods of the invention comprise obtaining at least a representative (e.g. a cross-section or circumferential surface) portion of stool voided by a patient, and performing an assay to detect in the sample the presence of cells or cellular debris shed from epithelial cells lining the colon that may be indicative of cancer or precancer. Most often, such cells will be derived from a polyp or a cancerous or precancerous lesion at a discrete location along the colon. For purposes of the present invention, a precancerous lesion comprises precancerous cells, and precancerous cells are cells that have a mutation that is associated with cancer and which renders such cells susceptible to becoming cancerous. As shown in Figure 1, a cross-sectional sample is a sample that contains at least a circumferential surface of the stool (or portion of a stool comprising an entire cross-sectional portion), as, for example, in a coronal section or a sagittal section. A sample comprising the surface layer of a stool (or of a cross-section of a stool) also contains at least a circumferential surface of the stool. Both cross-sections and circumferential surfaces comprise longitudinal stripes of sloughed colonic epithelium, and are therefore representative samples.

In a preferred embodiment, methods of the invention comprise the steps of obtaining at least a circumferential surface or cross-sectional portion of a stool voided by a patient, and performing an assay to detect debris indicative of disease. For example, such debris may comprise a clonal subpopulation of cells having one or more mutations (for purposes of the present application, a mutation is a deletion, substitution, addition, modification, intercalation or rearrangement of DNA). Preferred methods of the invention comprise detection of characteristics of such transformed cells, including, for example, mutations, proteins expressed uniquely or in altered

- 4 -

amounts in transformed cells, and blood. Particularly preferred methods of the invention comprise obtaining at least a circumferential surface or cross-sectional portion of a stool sample, and performing an assay to detect DNA characteristics indicative of the presence of a clonal subpopulation of cells in the sample. The clonal
5 subpopulation may be, for example, a subpopulation of cancerous or precancerous cells, having a mutation in, for example, a p53 tumor suppressor gene. Clonal subpopulations of cells detected by methods according to the invention are often characterized by a massive loss of DNA, resulting in a loss of heterozygosity that renders ineffective the gene or genes encompassed by the deletion. Alternative
10 methods of the invention comprise performing an assay to detect the presence of an infectious disease, for example debris from a microorganism.

Methods of the invention also comprise obtaining a representative (*i.e.*, cross-sectional or circumferential) sample of stool and homogenizing the stool in a buffer, such as a buffer comprising a detergent and a proteinase and optionally a DNase
15 inhibitor.

In methods according to the invention, an assay performed on at least a circumferential surface or cross-sectional portion of stool may be an assay to detect a disease when a diagnostic indicator of the disease is incorporated into stool. Methods of the invention are useful to detect cellular debris shed from the epithelial lining of the
20 colon, and from other tissue sources that shed cellular material into the gastrointestinal tract and preferably into the colon. Shed material may be from colonic epithelial cells, pancreatic cells, or may be indicative of infection (e.g. bacterial or viral nucleic acids or proteins). In methods according to the invention, an assay performed on at least a circumferential surface or cross-sectional portion of stool may be an assay to detect the
25 presence of elevated levels of carcinoembryonic antigen shed from cells lining the colon. Such an assay may also comprise detecting the presence of occult blood. However, methods of the invention preferably comprise an assay wherein the sample is exposed to an antibody that specifically binds to a molecule characteristic of cellular debris shed from cells comprising a subpopulation of cells having a mutation that is
30 potentially associated with cancer.

- 5 -

Methods of the invention are especially and most preferably useful for detecting DNA characteristics indicative of a subpopulation of transformed cells in a representative stool sample. The DNA characteristics may be, for example, mutations, including loss of heterozygosity, microsatellite instability, and others. An assay for

5 DNA characteristics in a method of the invention may comprise the step of determining whether a difference exists in a number X of a first allele known or suspected to be mutated in a subpopulation of cells in a representative stool sample, and a number Y of an allele known or suspected not to be mutated in the sample, a statistically-significant difference being indicative of a mutation and the possible presence of cancer in a

10 subpopulation of cells in the sample. In an embodiment of the invention, the difference between a number of a tumor suppressor gene and a number of a non-cancer-associated gene are compared, a statistically-significant difference in the numbers being indicative of a mutation in the tumor suppressor gene.

Assays useful in the practice of methods according to the invention also include

15 an assay to detect the presence of a deletion or other mutation in a region encompassing a polymorphic nucleotide. In such an assay, a number of a polymorphic nucleotide present at maternal and paternal alleles, wherein the patient is heterozygous for the polymorphic nucleotide, is determined. A statistically significant difference between a number of a polymorphic nucleotide in a maternal allele and a

20 paternal allele is indicative of the presence of a deletion in one of the two alleles. Another useful assay is one to detect an infectious disease.

Methods of the invention typically include, following sample preparation and an assay for characteristics of cells or cellular debris, a visual examination of the colon in order to determine if a polyp or other lesion is, in fact, present. Finally, surgical

25 resection of abnormal tissue may be done in order to prevent the spread of cancerous or precancerous tissue.

Accordingly, methods of the invention provide means for screening for the presence of a cancerous or precancerous subpopulation of cells in a heterogeneous sample, such as a stool sample. Methods of the invention reduce morbidity and

30 mortality associated with lesions of the colonic epithelium. Moreover, methods of the invention comprise more accurate screening methods than are currently available in

- 6 -

the art, because current methods take advantage of the observation that cancerous or precancerous cells shed debris only onto or into part of the surface of the forming stool. The present methods reliably assay over the entire circumference of the stool, thereby increasing the likelihood of detecting an abnormality if one exists. Further aspects and advantages of the invention are contained in the following detailed description thereof.

DESCRIPTION OF THE DRAWINGS

Figure 1 is a diagram of a cylinder which represents a formed stool and shows various cross-sections which will contain material from the entire circumference of a stool. The Section labeled "A" is a typical coronal section and the section labeled "B" is a typical sagittal section. The strip labeled "C" represents material shed from cancerous tissue which is deposited in a longitudinal stripe.

Figure 2 is a schematic diagram of a receptacle for containing a stool sample.

Figure 3 is a schematic diagram of a multi-orifice impedance counter; wherein reference numeral 1 indicates the direction of flow through the column; reference numeral 2 indicates a plunger means for forcing material downward in the column; reference numerals 3 and 4 are different-sized hybridization beads; reference numeral 5 is an optional filter for extracting unwanted particles; reference numeral 6 indicates an array of orifices for measuring differential impedance; and reference numeral 7 is a collection chamber.

Figure 4 is a diagram showing primers useful for the detection of single base polymorphisms.

DETAILED DESCRIPTION OF THE INVENTION

Methods according to the present invention are useful for the preparation of stool samples that will reproducibly contain cells or cellular debris shed from a clonal population of cancerous or precancerous cells, if such a population is present at any site along the colon of a patient. These samples are then used to perform assays to detect characteristics indicative of cancer in a highly-reproducible and accurate way. In preferred methods of the invention, a sample representative of the surface of a stool is obtained. Such methods provide an improvement over the art inasmuch as they teach removing the portion of stool onto which the colonic epithelium sheds debris.

- 7 -

Two preferred methods for obtaining such a portion are cross-sectioning and extracting the surface of the stool. In a more preferred method, the surface of a cross-section is obtained. Without the recognition that at least a cross-sectional sample must be obtained, there is no means for reproducibly obtaining a sample that will contain a cancerous or precancerous subpopulation of cells, if one exists.

Methods described in the art do not recognize that, unlike infection by parasites, bacteria and viruses, characteristics indicative of the presence of colon cancer, especially early stage colon cancer, are found only in a specific portion of voided stool. If the sampled portion of stool does not include the portion that happens to contain cells and cellular debris shed from early-stage cancer tissue, the diagnostic assay necessarily will fail to detect the characteristics indicative of the presence of colorectal cancer in a reliable manner even if homogenized, i.e., will produce a false-negative result. Methods of the invention are also useful to detect the presence of bacteria, virus or parasites, as they will also be present on the surface of formed stool.

Sloughed cells from, for example, a polyp forming in the epithelial lining of the colon, or on early stage cancerous lesions, are sloughed onto only the portion of the forming stool that comes into contact with the polyp or lesion. Accordingly, in early stage disease, only a small portion of the surface layer of the forming stool will contain sloughed cells, and if that portion happens not to be taken as part of the sample, an assay for indicia of colon cancer necessarily will produce a false-negative result. A brief review of the anatomy and physiology of the colon will aid in an understanding of this phenomenon.

A typical adult colon is approximately six feet in length, with a diameter of about two to three inches. Numerous bends and folds are present throughout its length. The colon removes water from liquid or semi-liquid waste material that enters the colon, and relatively solid stool begins to form in the proximal third of the colon. Epithelial cells line the lumen of the colon, and the luminal surface is organized into microscopic crypts. Colorectal epithelial cells are replaced every four to five days. The epithelial cells divide rapidly at the base of the crypts and migrate to the apices, where cells appear to undergo apoptosis (programmed cell death), and cellular debris is shed into the lumen. The lining of the colorectal lumen is elastic and the diameter of the lumen is

- 8 -

determined by the volume of stool that is passing through the colon at any given time. As a result, the surface of the forming stool passing through the colon is in direct contact with the epithelial lining of the lumen. Shed epithelial cells (which may or may not have undergone apoptosis) and cellular debris therefore are incorporated onto the
5 surface of stool as it passes through the colon.

Cells and cellular debris from colorectal epithelial cancers therefore also are shed onto forming stool. Most colorectal cancers develop in regions of the colon where stool is relatively solid, indeed approximately one third of such cancers develop in the rectum. Markers indicative of the presence of cancer, including cells, cellular debris,
10 DNA, blood, and carcinoembryonic antigen, are shed onto the portion of the forming stool that contacts the cancerous tissue as the stool passes through the colon. Since the stool is relatively solid, these markers will remain on or near the surface of the stool where they were deposited and will not be homogeneously dispersed throughout the stool. As stool passes over a cancerous or precancerous growth, material from the
15 growth will be deposited along the stool, but only on the part of the stool circumference that comes into direct contact with the cancerous or precancerous tissue comprising the lesion. Stool voided by a patient with colorectal cancer or precancer is therefore characterized by a longitudinal "stripe" of diagnostically relevant material derived from the cancerous or precancerous tissue.

20 A sample that does not include material from the entire circumference of a stool voided by a patient with colorectal cancer or precancer will not reproducibly contain material derived from the cancerous or precancerous tissue. Currently, random, non-cross-sectional samples ("smears") of voided stool are analyzed in clinical settings. In these, sloughed cancerous or precancerous cells and cellular debris have no
25 possibility of detection unless the sample happens by chance to contain the portion of stool which made contact with the region of the colon from which cells were sloughed.

Furthermore, cancers typically develop by clonal expansion of a single mutant cell, and in the early stages of the disease, i.e., when surgical removal is an effective cure, the cancerous lesion will be very small and may lie on a small arc of the
30 circumference of the colon. Material derived from such an early stage cancer therefore will be shed onto or into stool in a very narrow stripe (labeled C in Figure 1).

- 9 -

Consequently, a sample that does not contain the entire circumference of a stool voided by a patient with early stage colorectal cancer or precancer only by chance will contain material indicative of the presence of the early stage cancerous or precancerous condition. However, early detection of colorectal cancer is very
5 important for effective surgical intervention. The present invention provides methods for reproducible early detection of characteristics indicative of the presence of cancer or precancer in a patient.

Similarly, the present invention provides methods for reproducibly detecting an indicator of the presence of an infectious agent, such as a virus, bacterium, parasite or
10 other microorganism, in stool from an infected patient. Methods of the invention are particularly useful for reproducibly detecting an infectious agent, or an indicator characteristic thereof, that is deposited onto narrow area of stool as it forms in the colon.

Analysis of at least a circumferential surface or cross-sectional sample of stool
15 ensures that a disease indicator, incorporated on a stool as it moves through the colon, will be present in a diagnostic assay sample. For example, analysis of at least a circumferential surface or cross-sectional sample of stool (including a whole stool), as shown in Figure 1, ensures that at least a portion of cells and cellular debris shed from any existing cancerous or pre-cancerous cells (even if shed from small early stage
20 cancerous or pre-cancerous tissue, e.g., small polyps) will be present in the portion of the stool sample to be analyzed. Indeed taking at least a circumferential surface or cross-section of the stool sample avoids the possibility of analyzing stool portions that will not contain sloughed cancerous or precancerous cells even when the patient has colorectal cancer or precancer.

25 Once a circumferential surface or cross-sectional sample of stool is obtained, it may be homogenized by known methods to distribute cells and cellular debris throughout the sample. An assay then is performed on the homogenate, or an extract of the homogenate, to detect the presence of cells and/or cellular debris in the sample. The assay may be any one or a combination of histological cellular assays, antibody
30 based immunoassays (or other formats) designed to detect the presence of a molecule characteristic of transformation such as a protein, or DNA-based assays for detecting

- 10 -

mutations or genetic characteristics indicative of colorectal cancer. Known assay protocols, those disclosed herein or in U.S. Patent No. 5,670,325, or assays hereafter developed may be used in the practice of the invention. Non-limiting examples of useful known assay protocols include those disclosed in U.S. Patent Nos. 5,137,806
5 (detection of sequences in selected DNA molecules), 5,348,855 (assay for nucleic acid sequences), 5,512,441 (detection of mutant alleles), 5,272,057 and 5,380,645 (RFLP analysis), 5,527,676 (detection of p53 gene sequences), 5,330,892 (detection of MCC gene sequences), 5,352,775 (detection of APC gene sequences), 5,532,108 (detection of DCC gene sequences), and in WO96/08514 (monoclonal antibodies against human
10 colon carcinoma-associated antigens), the disclosures of each of which are incorporated by reference herein. Alternatively, or additionally, an assay for fecal occult blood may be performed as reported in U.S. Patent Nos. 4,333,734 and 5,196,167, incorporated by reference herein. Assays useful in the context of the present invention also include an assay for carcinoembryonic antigen as reported in
15 U.S. Patent No. 5,380,647, incorporated by reference herein. In addition, assays described in USSN 08/808,763; 08/815,576; 08/876,857; 08/876,638; 08/877,333; and 60/063,219; incorporated by reference herein, are also useful in the context of the present invention. Finally, the sample may be prepared, as reported in U.S. Patent No. 4,857,300, incorporated by reference herein, for histological examination to detect
20 characteristics indicative of the presence of cancerous or precancerous cells.

The purpose of any assay protocol used in connection with obtaining at least a cross-sectional sample is to identify candidates for subsequent invasive diagnostic procedure such as colonoscopy or sigmoidoscopy. The assay accordingly need not definitively detect the presence of a cancerous or precancerous lesion, although false
25 negatives obviously are to be avoided. The goal of the test protocol is not to determine whether in the vast quantities of cell debris in the sample there are a few cells which bear a mutation commonly associated with early stage transformation, but rather whether the sample contains debris indicative of clonal expansion of a mutant cellular subpopulation. Maximum benefit will come from an assay designed to detect the likely
30 presence of a clonally expanded cell population, i.e., of transformed colonic epithelial cells which comprise the cancerous or precancerous lesions. Assays having the ability

- 11 -

to detect cancer or precancer in early stages are preferred in methods of the invention. Assays using polymerase chain reaction (PCR), restriction fragment length polymorphism (RFLP) or other methods for nucleic acid analysis may be used to detect known DNA characteristics indicative of the presence of colorectal cancer or
5 precancer. More precise methods for quantitative detection of cellular debris, such as DNA fragments or segments, may be used to analyze cross-sectional samples according to methods described herein.

A preferred assay interrogates the sample for DNA characteristics indicative of the development of cancer or precancer. However, assays for use with methods of the
10 invention may detect any abnormal cellular debris shed from clinically-relevant transformed tissue. Thus, in accordance with a preferred aspect of the invention, an assay is used to detect the presence of characteristics of cells which have experienced loss of heterozygosity, microsatellite instability or other mutation.

The following examples provide details of methods according to the invention.
15 However, numerous additional aspects of the invention, especially in terms of assays to be performed, will become apparent upon consideration of the following detailed description thereof.

Example 1

Preparation of a Stool Sample

20 A sample is prepared such that it contains at least a circumferential surface or cross-sectional portion of a stool voided by a patient.

Stool is voided into a receptacle that is preferably small enough to be transported to a testing facility. The receptacle may be fitted to a conventional toilet such that the receptacle accepts stool voided in a conventional manner. The
25 receptacle may comprise a mesh or screen of sufficient size and placement such that stool is retained while urine is allowed to pass through the mesh or screen and into the toilet. The receptacle additionally may comprise means for removing a circumferential surface or cross-sectional portion from the stool. Moreover, the receptacle may comprise means for introducing homogenization buffer or one or more preservatives,
30 such as alcohol, a solution of high salt concentration, antibiotics, and chaotropic salts in order to neutralize bacteria present in the stool sample. The homogenization buffer

- 12 -

may be a physiologically compatible buffer such as phosphate buffered saline, and may comprise salt such as 20-100 mM NaCl or KCl. The homogenization buffer may also comprise a detergent, such as 1-10% SDS or triton, and/or a proteinase, such as proteinase K. The buffer may also contain inhibitors of DNA and RNA degrading enzymes.

The receptacle, whether adapted to fit a toilet or simply adapted for receiving the voided stool sample, should include sealing means sufficient to contain the voided stool sample and any solution added thereto and to prevent the emanation of odors. An exemplary receptacle is shown in Figure 2. A shown in that figure, the receptacle has a support frame 1 which is placed directly over the toilet bowl 2. The support frame 1 has attached thereto an articulating cover 3 which may be placed in a raised position, as shown in figure 2, for depositing of sample or a closed position (not shown) for sealing voided stool within the receptacle. The support frame 1 additionally has a central opening 4 traversing from a top surface 5 through to a bottom surface 6 of the support frame 1. The bottom surface 6 directly communicates with a top surface 7 of the toilet 2. Extending from the bottom surface 6 of the support frame 1 is a means 8 for capturing voided stool. Means 8 may be fixedly attached to the support frame 1 or may be removably attached for removal subsequent to deposition of stool. Means 8 may comprise a further means for removing at least a circumferential surface or cross-sectional portion from the voided stool. A preferable sample size is at least 5-10 g or at least 5-10 ml. A means to assess the presence of a minimal sample size may comprise a physical diagram indicating the minimal sample size. Alternatively a means to assess the presence of a minimal sample size may comprise the displacement of a liquid or of a mechanical device to a minimal level upon deposit of the stool sample.

In a preferred embodiment, a cross-sectional portion is removed from the voided stool by making one or more sagittal or coronal section through the stool, as shown in Figure 1. The removed portion comprises material from an entire circumferential surface of the stool. Alternatively, a circumferential surface of a stool is prepared by taking the entire surface layer of the stool, or the entire surface layer of a cross-sectional portion of the stool. In a preferred embodiment, the surface layer is approximately 0.001 to 0.1 mm thick. In another preferred embodiment, the surface

- 13 -

layer is approximately 0.1 to 20 mm thick. In a more preferred embodiment, the surface layer is approximately 5 mm thick. In a most preferred embodiment, the surface layer is representative of material deposited on the forming stool, and reproducibly contains markers or indicators, such as shed cellular material, of a disease that is present in the

5 patient. In one embodiment, a stool surface layer is obtained by cutting or scraping off the surface of the stool or cross-sectional portion of the stool. In a preferred embodiment, a stool surface layer is prepared using a cylindrical cutting device (for example a rigid loop) that has a diameter that is smaller than the diameter of the stool or cross-sectional portion of the stool. The surface layer is obtained when the stool

10 sample is forced through the cutting device. In a preferred embodiment, the diameter of the cutting device is adjustable. Alternatively, a circumferential surface of a stool is obtained by contacting the surface of a stool or cross-sectional portion of a stool with a sampling device. In one embodiment, the sampling device is first wrapped around the stool sample and subsequently removed or peeled off. The layer of stool that is peeled

15 off with the sampling device is then processed as described herein. Alternatively, the stool sample is rolled along the surface of the sampling device, and the layer of stool that adheres to the surface is then processed as described herein. Alternatively, a diagnostic assay is performed directly on the layer of material on the sampling device. It is important that the entire circumference of the stool is sampled. In a preferred

20 embodiment, the sampling device has a surface that stool adheres to. In a more preferred embodiment, the sampling device surface preferentially binds diagnostic markers or indicators of a disease (for example nucleic acid or protein molecules), and thereby removes them from the surface of the stool sample. In one embodiment, the sampling device is an adherent paper or membrane. In a preferred embodiment, a

25 stool is frozen prior to processing. In one embodiment the stool is frozen to between -5 and -30 °C. In a preferred embodiment the cross-sectional portion or circumference of the stool is obtained from the frozen or partially thawed stool. In an alternative embodiment, a whole stool may be used for diagnostic assays. The stool or portion of stool that is obtained contains sufficient material to allow subsequent diagnostic assays

30 to be performed.

- 14 -

Once obtained, the entire circumference or cross-sectional sample of stool is homogenized in an appropriate buffer, such as phosphate buffered saline.

Homogenization means and materials for homogenization are generally known in the art. Thus, particular homogenization methods may be selected by the skilled artisan

5 and may depend upon the assay to be employed. The buffer may contain detergent, salt, proteinase, inhibitors of DNA and RNA degrading enzymes. The composition of the buffer will depend on the type of assay to be performed. If a fecal occult blood assay is to be performed, the buffer may contain chemical compounds which react with blood to produce a color, the intensity of which can be measured. Buffers useful for
10 detecting the presence of fecal occult blood are known in the art. If a test is to be performed for the presence of a particular protein, the buffer should not contain a proteinase capable of degrading such tumor marking antigens.

DNA or RNA may be isolated from the homogenate using methods known in the art. Subsequent tests may be performed on the isolated DNA and RNA.

15 **Example 2**

Exemplary Enumerative methods for Detection of Colorectal Cancer or Precancer in stool samples

DNA characteristics associated with the presence of colorectal cancer or a precancerous lesion may be detected in stool samples prepared according to the
20 invention, using, for example, the methods described in the following sections. A careful endoscopic examination preferably is performed on positive Individuals, followed by early surgical excision of any diseased tissue.

A. Reference-Target

Methods of the invention are used to prepare a stool sample followed by
25 detection of a deletion or other mutation in the p53 tumor suppressor gene. The p53 gene is a good choice because a loss of heterozygosity in p53 is often associated with colorectal cancer. An mRNA sequence corresponding to the DNA coding region for p53 is reported as GenBank Accession No. M92424. At least a cross-section of a voided stool sample is obtained and prepared according to methods of the invention as
30 described immediately above. The sample need not be further processed for analysis. However, DNA or RNA may optionally be isolated from the sample according to

- 15 -

methods known in the art. See, Smith-Ravin, *et al.*, *Gut*, 36: 81-86 (1995), incorporated by reference herein.

5 Nucleic acids may be sheared or cut into small fragments by, for example, restriction digestion. The size of nucleic acid fragments produced is not critical, subject to the limitations described below. A target allele that is suspected of being mutated (p53 in this example) and a reference allele are chosen. A reference allele may be any allele known normally not to be mutated in colon cancer.

10 Either portions of a coding strand or its complement may be detected. For exemplification, detection of the coding strand of p53 and reference allele are described herein. Complement to both p53 and reference allele are removed by hybridization to anti-complement oligonucleotide probes (isolation probes) and subsequent removal of duplex formed thereby. Methods for removal of complement strands from a mixture of single-stranded oligonucleotides are known and include techniques such as affinity chromatography. Upon converting double-stranded DNA to
15 single-stranded DNA [See, e.g., Sambrook, *et al.*, *Molecular Cloning, A Laboratory Manual* (1989) incorporated by reference herein], sample is passed through an affinity column packed with bound isolation probe that is complementary to the sequence to be isolated away from the sample. Conventional column chromatography is appropriate for isolation of complement. An affinity column packed with sepharose or other
20 appropriate materials with attached complementary nucleotides may be used to isolate complement DNA in the column, while allowing DNA to be analyzed to pass through the column. See Sambrook, *Supra*. As an alternative, isolation beads may be used to exclude complement as discussed in detail below.

25 After removal of complement strands, first oligonucleotide probes which hybridize to at least a portion of the p53 allele and second oligonucleotide probes that hybridize to at least a portion of the reference allele are obtained. The probes are labeled with a detectable label, such as fluorescein or with detectable particles. Distinct labels for the probes are preferred. However, identical labels may be used if, for example, sample is assayed in two separate aliquots. Probes may be labeled with
30 identical or with distinct labels. However, distinct labels are preferred.

- 16 -

Labeled probes then are exposed to sample under hybridization conditions. Such conditions are well-known in the art. See, e.g., Wallace, et al., *Nucleic Acids Res.*, 6:3543-3557 (1979), incorporated by reference herein. First and Second oligonucleotide probes that are distinctly labeled (i.e. with different radioactive
5 isotopes, fluorescent means, or with beads of different size, See *infra*) are applied to a single aliquot of sample. After exposure of the probes to sample under hybridization conditions, sample is washed to remove any unhybridized probe. Thereafter, hybridized probes are detected separately for p53 hybrids and reference allele hybrids. Standards may be used to establish background and to equilibrate results. Also, if
10 differential fluorescent labels are used, the number of probes may be determined by counting differential fluorescent events in a sample that has been diluted sufficiently to enable detection of single fluorescent events in the sample. Duplicate samples may be analyzed in order to confirm the accuracy of results obtained.

If there is a statistically-significant difference between the amount of p53
15 detected and the amount of the reference allele detected, it may be assumed that a mutation has occurred in p53 so as to alter its sequence and prevent hybridization of the probe, or that at least a portion of the region of the genome containing p53 has been lost in a subpopulation of cells shed from the colon. The patient therefore may be at risk for developing or may have developed colon cancer. Statistical significance may
20 be determined by any known method. See, e.g., Steel, *et al.*, Principles and Procedures of Statistics: A Biometrical Approach (McGraw, Hill, 1980). Statistical methods are also outlined in co-owned U.S. Patent No. 5,670,325.

The determination of a p53 mutation allows a clinician to recommend further treatment, such as endoscopy procedures, in order to further diagnose and, if
25 necessary, treat the patient's condition. The following examples illustrate methods that allow direct quantification of hybridization events.

1. Method for Quantitation of Target and Reference Polynucleotides

Enhanced quantification of binding events between hybridization probes and
30 target or reference is accomplished by coupling hybridization probes to particles, such as beads (hybridization beads). In order to obtain a precise quantitative measure of

- 17 -

the amount of a polynucleotide in a sample, hybridization beads are constructed such that each bead has attached thereto a single oligonucleotide probe.

a. *Method for Preparation of Probe-Bead Combinations*

A single probe is attached to a bead by incubating a large excess of
5 hybridization beads with oligonucleotide probes of a given type (*i.e.*, either first or second oligonucleotide probes). Coupling of probe to bead is accomplished using an affinity-binding pair. For example, beads may be coated with avidin or streptavidin and probes may be labeled with biotin to effect attachment of the probe to the bead. The mixture of beads and probes is agitated such that 100% of the probes are bound to a
10 bead. The mixture is then exposed to a matrix, such as an affinity column or a membrane coated with oligonucleotides that are complementary to the probe. Only beads that have an attached probe will adhere to the matrix, the rest being washed away. Beads with coupled probe are then released from the matrix by melting hybridizations between probe and complement. Multiple exposures to the matrix and
15 pre-washing of the column reduces non-specific binding. Moreover, naked beads (*i.e.*, without attached probe) may be exposed to the matrix to determine a background number of beads that can be expected to attach to the matrix in the absence of probe.

By using a vast excess of beads relative to probe as described above, the vast majority of recovered beads will have only one attached probe. For example, if a
20 mixture has a ratio of 1 probe to 1000 beads, it is expected that only about 1 bead in a million will have two attached probes and even less than one bead in a million will have more than two attached probes. Accordingly, hybridization beads are provided in an effective 1:1 ratio with probe which allows for precise quantification of target and reference polynucleotide as described below.

25 For each assay described below, two distinct hybridization beads are used. A first hybridization bead has attached thereto a single first oligonucleotide probe that is complementary to at least a portion of a target polynucleotide (*e.g.*, a p53 allele). A second hybridization bead, of a size distinct from the first hybridization bead, has attached thereto a single second oligonucleotide probe that is complementary to at
30 least a portion of a reference polynucleotide (*i.e.*, one that is known or suspected not to be mutated in the sample).

b. *Use of Beads to Quantify Target and Reference Polynucleotides*

5 DNA is melted (denatured to form single-stranded DNA) by well-known methods See, e.g., Gyllenstein, *et al.*, in *Recombinant DNA Methodology II*, 565-578 (Wu, ed., 1995), incorporated by reference herein. One may detect either a coding strand or its complement in order to quantify target and/or reference polynucleotide. For purposes of illustration, the present example assumes detection of the coding strand.

10

2. *Removal of Complement*

Single-stranded complement of the target polynucleotide (e.g., p53) and reference polynucleotide are removed from the sample by binding to oligonucleotide probes that are complementary to target or reference complement. Such probes, referred to herein as isolation probes, are attached to isolation beads prior to their introduction into the sample. The beads may be magnetized. Thus, when magnetized isolation beads [with attached isolation probe(s)] are introduced into the sample, the attached isolation probes hybridize to complement of target or reference (or vice versa). Isolation beads preferably are introduced in vast excess in order to saturate complement binding. Once hybridization is complete, a magnetic field is applied to the sample to draw the magnetized isolation beads (both with and without hybridized complement) out of the sample. Assuming that a sufficient quantity of isolation beads are introduced into the sample, removal of the isolation beads effectively removes all target and reference complement from the sample.

25 In an alternative method for complement removal, an excess of oligonucleotide probe labeled with biotin is exposed to the melted or dehybridized (single stranded) sample under hybridization conditions. Once hybridization is complete, the sample is exposed to a column containing immobilized avidin. The biotin-labeled probe, whether free or hybridized to complement, is bound by avidin on the column. The remainder of the DNA, including target and reference coding strands to be detected, pass through
30 the column. In contrast to the description of hybridization beads above, beads for

- 19 -

removal of complement may each comprise multiple complementary oligonucleotide probes.

3. *Quantitation of Target and Reference*

Two sets of hybridization beads are prepared as described above. Each
5 member of a first set of hybridization beads (all of which are identical to each other)
has attached thereto a single oligonucleotide probe that is complementary to at least a
portion of the target polynucleotide, i.e., the portion of the genome which is altered in
the cells of a cancerous lesion. Each member of a second set of identical hybridization
beads (all of which are identical to each other but not to the first set) has attached
10 thereto a single oligonucleotide probe that is complementary to at least a portion of the
reference polynucleotide, i.e., a portion of the genome which is not likely to be altered
in malignant cells. Members of the second set of hybridization beads are of a size or
color distinct from that of members of the first set of hybridization beads. First and
second hybridization beads may also be distinguished on the basis of other
15 characteristics. For example, beads may have fluorescent markers that are
distinguished by their fluorescence wavelength. Beads with distinct electrochemical
charges also may be used. The precise modality used for distinguishing beads is not
essential as long as it is possible to distinguish between first and second probe on the
basis of distinctions between attached first and second beads.

20 Both sets of hybridization beads are exposed to the sample under hybridization
conditions thereby allowing hybridization to reference and target. The sample then is
washed to remove unhybridized bead/probe combinations. Unhybridized bead/probe
combinations are removed by, for example, passing the sample through a column lined
of immobilized DNA complementary to the probe sequence. Thus, any unhybridized
25 bead/probe combinations are retained on the column while duplex passes through.
Subsequently, the sample is exposed to means for differentially counting hybridization
beads in order to quantify first and second hybridization probes which have formed
duplexes. The numbers obtained provide a precise estimate of the number of copies of
the reference and target polynucleotide in the population because differential counting
30 means count individual beads. One bead is equal to one probe which, in turn, signifies
one copy of the nucleic acid being measured.

- 20 -

An example of a differential counting means is an impedance measuring device, such as a Coulter counter (Coulter Electronics, Inc., Miami, Florida). Sample is passed through the device which differentially detects the two types of hybridization beads by measuring their differential impedance of an electric current. Alternatively, the device
5 may measure fluorescence, color, or other parameters. In order to increase the speed of the assay, a multi-orifice device may be used. A multi-orifice impedance counter is shown schematically in figure 2. A multi-orifice array is placed at one end of a column filled with an electrically-conductive fluid, such as saline. Hybridization beads with either hybridized target or reference segments are inserted at an opposite end of the
10 column. Each orifice is large enough to accommodate only one hybridization bead at a time and sufficiently wide to allow reliable impedance measurements. A voltage is set across each orifice. Each hybridization bead (which is non-conducting), as it passes through one of the orifices, displaces a volume of saline, thus creating an impedance that is proportional to its size. This, in turn, creates a measurable decrease in current
15 that is directly correlated with the size of the bead. By compiling the number of each of the two distinct impedance events, a precise estimate of the number of hybridization beads and, therefore, the number of probes of each type in the population may be obtained.

Upon quantitative measurement of first and second hybridization beads, the data
20 may be analyzed to determine whether any difference between the amounts of first and second hybridization beads is statistically significant. A reduction in the amount of target relative to the reference is indicative of a mutation in or deletion of the target allele in a subpopulation of cells in the sample. Where the p53 gene is the target allele, such a mutation is indicative of a cancerous or precancerous condition. A
25 clinician may use such results as a basis for prescribing additional treatment, such as endoscopy and polypectomy procedures.

B. Detection of Mutations in Single-base Polymorphisms

The basic method described above may also be applied to detect a loss of heterozygosity or other mutation at a single base polymorphic site between maternal
30 and paternal alleles. Such detection is typically an indication of a larger deletion or other mutation. However, a mutation at a single polymorphic nucleotide may be all that

- 21 -

is necessary to inhibit gene function in one of the two alleles. A mutation in a single-base polymorphic region may be difficult to detect due to a recently-discovered phenomenon called complementary reduplication. In complementary reduplication, the loss of one of two alleles at a particular locus results in "reduplication" of the surviving allele. Reduplication usually takes place on the chromosome containing the surviving allele and involves the production of one or more copies of the surviving allele in close proximity on the chromosome to the position of the surviving allele. In the case of a locus that displays one or more single-base allelic polymorphisms (*i.e.*, heterozygosity at the locus is determined by virtue of one or more single-base differences in one or more regions of the locus), complementary reduplication results in the insertion on the chromosome containing the surviving allele of a duplicate of the sequence corresponding to that which was deleted. Even under the most stringent hybridization conditions, some of a probe directed against the deleted sequence will bind to the reduplicated sequence at a locus of a single-base polymorphism. Accordingly, in such circumstances, the deletion may not be detected because any true difference in the number of probes binding to the polymorphic site (*i.e.*, the allelic region encompassing the single-base polymorphism) may be obscured by an increase resulting from the other allele's reduplicated region.

The problems associated with complementary reduplication, and with non-specific probe binding generally, are alleviated by the practice of the methods described herein. Such methods allow detection of a deletion in one of two alleles present at a specific locus in a subpopulation of cells contained in a biological sample. Numerous alleles, including tumor suppressor alleles, contain single polymorphic nucleotides in the context of a constant nucleic acid region. Individuals normally may be either homozygous or heterozygous for the polymorphic nucleotide. Since numerous single-base polymorphic nucleotide sites exist in most alleles, the probability that a given individual is heterozygous at least one of the single-base polymorphism sites is high. A statistically-significant reduction in one of the two nucleotides at a single-base polymorphic site (at which the individual is heterozygous) may be used as a marker for a deletion in the allele encompassing that site.

- 22 -

Genomic regions containing known single-base polymorphisms may be identified by reference to a nucleotide database, such as GenBank, EMBL, or any other appropriate database. The existence of polymorphisms may be determined by methods taught herein, gel electrophoresis or by other standard methods. For purposes of the invention, a single-base polymorphism is intended to be a single polymorphic nucleotide adjacent to a non-polymorphic region of the allele regardless of whether the single polymorphic nucleotide forms part of a larger polymorphic site (i.e. the single-base polymorphism may be the terminal nucleotide of a larger, polynucleotide polymorphism). For cancer detection, the regions considered are regions in which loss of heterozygosity is prevalent, such as regions containing tumor suppressor genes. A given individual may be homozygous or heterozygous for the polymorphic nucleotide in any identified single-base polymorphic region. Accordingly, if a number of single-base polymorphic regions are identified, the probability increases that at least one heterozygous single-base polymorphic region is found in a sample.

Once single-base polymorphic sites are identified, a DNA sample is obtained from a patient, e.g., from blood cells, to determine which of those sites is heterozygous in normal (i.e., non-cancerous or non pre-cancerous) cells for that individual. Then, a stool sample is prepared as described above. Double stranded DNA in the sample is converted to single-stranded DNA. Then, either the coding strand or the anti-coding strand for both alleles is removed from the sample. As will be evident from the following discussion, methods disclosed herein are indifferent as to whether coding strand or anti-coding strand is tested.

An oligonucleotide probe is constructed that is complementary to a portion of the region of single-base polymorphism, said portion ending at the nucleotide that is immediately 3' to the polymorphic nucleotide, regardless of whether the 5'-3' (coding) strand or the 3'-5' (anticoding) strand is used as a template. Figure 3 shows four possible probes that are immediately 3' to the polymorphic nucleotide for each of four possible template strands as described above (the Sequences in Figure 3 are hypothetical and are not intended to represent any actual sequence). The sequence labeled M1 is SEQ ID NO:1; the sequence labeled M2 is SEQ ID NO:2; the sequence

- 23 -

labeled M3 is SEQ ID NO: 3; the sequence labeled M4 is SEQ ID NO:4; the sequence labeled F1 is SEQ ID NO: 5; the sequence labeled F2 is SEQ ID NO: 6; the sequence labeled F3 is SEQ ID NO: 7; and the sequence labeled F4 is SEQ ID NO: 8. While either strand may be used as a template for probe binding to determine heterozygosity and/or the loss thereof, the sequence of the probe that is hybridized to the template will be different depending upon the strand used. Probes may be of any length that allows efficient and specific hybridization. Figure 3 merely illustrates four hypothetical probes that are useful for hybridization to the hypothetical sequence shown. The length of probe sequences may be determined as appropriate for each genomic region that is analyzed. A preferable length is between about 10 and about 100 nucleotides. The size of the probe will also depend upon the size of the region surrounding the single-base polymorphism (*i.e.*, the region 5' or 3' to the next adjacent polymorphism, if any). Details concerning the construction and hybridization of oligonucleotide probes are known in the art.

Unique probes for each polymorphic region will hybridize to regions of both maternal and paternal alleles up to, but not including, the polymorphic nucleotide, which, in a heterozygote, will be different in the maternal and paternal alleles. Figure 3 shows only a small portion of the region surrounding the polymorphic nucleotide. The alleles shown in Figure 3 are heterozygous at the polymorphic site.

Probe is hybridized to its specific template DNA by standard methods. The sample may optionally be washed to remove unhybridized probe. To determine whether each target region bound by a probe is heterozygous or homozygous at the polymorphic nucleotide, a modification of the dideoxy chain termination method as reported in Sanger, *Proc. Nat'l Acad. Sci. (USA)*, 74: 5463-5467 (1977), incorporated by reference herein, is used. The method involves using at least two of the four common 2', 3'-dideoxy nucleoside triphosphates (ddATP, ddCTP, ddGTP, and ddTTP). A different detectable label is attached to each dideoxy nucleoside triphosphate (ddNTP) according to methods known in the art. Differentially-labeled ddNTPs are available commercially, for example, from Perkin Elmer Corporation (Cat. No. 401456).

At least two labeled ddNTPs then are exposed to each sample having probe hybridized to maternal and paternal alleles as described above. The choice of which two ddNTPs

- 24 -

are used will depend upon the nucleotides at the heterozygous polymorphic site. Any 3' modified nucleoside triphosphate may be used in the method as long as the 3' modification prevents binding of an additional 3' nucleotide (i.e. probe extension) and does not inhibit binding of the modified nucleotide to the 3' end of the probe. A DNA
5 polymerase, such as Sequenase™ (Perkin-Elmer), is added to the sample mixture. Using the allelic strands as primer, the polymerase will add one ddNTP to the 3' end of the probe, the incorporated ddNTP will be complementary to the nucleotide that exists at the single-base polymorphic site. Because the ddNTPs have no 3' hydroxyl, further elongation of the hybridized probe will not occur. After completion, the sample is
10 washed to remove excess ddNTPs. Label is then counted in each sample. The presence of two differentially-labeled ddNTPs in a sample is indicative of heterozygosity at the polymorphic site.

It is not necessary to determine the amount of each label present in the sample in order to establish heterozygosity or homozygosity. For Example, differentially-
15 labeled deoxynucleoside triphosphates may be used for a determination of heterozygosity or homozygosity. The mere fact that two different labeled dideoxy nucleotides are incorporated into the probe means that the single-base polymorphic site being analyzed is heterozygous. However, determination of sites at which a patient is polymorphic is useful in order to establish a baseline of polymorphisms which
20 may be used in future tests to detect changes in polymorphic sites which may be indicative of cancer. The existence of polymorphisms may be determined by methods taught herein, by gel electrophoresis or by other standard methods.

In the case in which heterozygosity exists at the polymorphic site, counting the amount of each of the two differentially-labeled ddNTPs allows a determination of
25 whether there is a loss of heterozygosity (i.e., a deletion) in a subpopulation of cells in the sample. In a normal (i.e., non-cancerous) sample containing cells that are heterozygous at the single-base polymorphic site, it is expected that the detected amount of each of the two ddNTPs added to the probe will be identical (within chosen limits of statistical significance). However, if a deletion has occurred in one of the two
30 alleles in a subpopulation of cells in the sample, there will be a statistically-significant difference between the amounts of each of the two alleles detected via the incorporated

- 25 -

(labeled) ddNTPs. The detection of such a difference is indicative of genomic instability within the sample. Such genomic instability indicates the possibility of cancerous or pre-cancerous cells in the sample.

In order to improve the ability to count alleles to which ddNTPs have attached accurately, ddNTPs are labeled with hybridization-type beads of different sizes as described above. Alleles with bound probe comprising a labeled ddNTP are counted as described above using a counting device, such as a Coulter counter. Also as described above, differential fluorescent labels or other counting means may be used to separately detect incorporated ddNTPs.

The detection of heterozygosity at single-base polymorphic sites and the detection of the loss of heterozygosity may be determined in separate steps. For example, probes may be hybridized immediately adjacent to but not including the nucleotide determined to be polymorphic as described above. The four ddNTPs may then be added to the sample, washed, and the presence or absence of each label may be detected. Detection of only one label indicates that the individual from whom the sample was obtained is homozygous at the site of the potential polymorphic nucleotide. Detection of two labels means that the individual is heterozygous. The heterozygous loci are recorded. As noted above, baseline determinations of heterozygosity may be done using standard methods. Once a baseline is established, future tests on that individual are performed exploiting the heterozygous loci in order to detect a loss of heterozygosity. For the detection of cancer, the heterozygous loci are typically chromosomal areas containing tumor suppressor genes, including p53, dcc, apc, and others. Using methods described herein, a "fingerprint" of heterozygous tumor suppressor loci may be constructed. Future deviation from the fingerprint (*i.e.*, deletions) provides valuable information as to the development of cancer.

A preferred use of the foregoing methods is in the detection of colon cancer. A representative stool sample is prepared as described above. Double-stranded DNA is converted to single-stranded DNA and complement of the strand to be detected is removed from the sample. The remaining single-stranded DNA is exposed to multiple copies of a probe designed on the basis of known single-base polymorphisms in a cancer-associated allele such that the probe hybridizes with a desired number of

- 26 -

nucleotides immediately adjacent the polymorphic nucleotide as described above. After hybridization is complete, the sample is washed and exposed to differentially-labeled ddNTPs and a DNA polymerase. The sample then is washed to remove unincorporated ddNTPs. The presence of any labeled ddNTPs is determined. If two
5 labels are detected, the individual from whom the sample is obtained is heterozygous at the polymorphic nucleotide. The heterozygosity of the allele and the probe sequence matching the site immediately adjacent to the polymorphic allele are noted for reference in future testing for the loss of heterozygosity. Alternatively, once the patient is determined to be heterozygous at a locus, an assay may be performed
10 immediately in the manner described above in order to determine an existing loss of heterozygosity in a subpopulation of cells in the sample.

C. Analysis of Microsatellite Instability

Microsatellites are di- or trinucleotide repeats found throughout the genome. A particular array of microsatellite repeats is often associated with a particular genomic
15 sequence and is stably inherited under normal conditions. Expansions of microsatellite copy number typically, called "microsatellite instability," are associated with defects in mismatch repair. Accordingly, changes in a microsatellite region indicate that the patient is at risk for a mutation in other genomic regions.

In order to detect microsatellite instability as an indicator of a mutation in a
20 cancer-associated gene, one must first identify a microsatellite region associated with the gene of interest. Such regions are typically identified on a database, such as GenBank, EMBL, and others. Once a wild-type microsatellite region associated with, for example, the p53 tumor suppressor gene, is identified, an oligonucleotide probe is constructed that spans the microsatellite region and the regions immediately 5' and
25 immediately 3' to the microsatellite region. The precise length of probes may be determined by the experimenter. Probes are constructed that hybridize to the microsatellite region, including portions extending 5' and 3', on both the maternal and paternal alleles with which the microsatellite is associated (e.g., p53).

An appropriate sample of body tissue or fluid is obtained and processed as
30 described herein. Double stranded DNA is denatured and an excess of maternal and

- 27 -

paternal probes, as described above, are introduced into the sample under hybridization conditions. The probes are detectably labeled as described above. Complement of the strands to be detected may optionally be removed by methods described above. The sample is then washed to remove unhybridized probe and the amount of hybridized probe is quantitatively detected.

Quantitative detection may be accomplished by any means described herein. For example, probes may be attached to hybridization beads such that probes that bind to maternal allele are attached to beads of one size and probes that bind to paternal allele are attached to beads of a second size that is distinguishable from beads of the first size. Beads with attached probe may be counted as described above.

The detection of a statistically-significant difference between the amount of probe binding to the maternal allele and the amount of probe binding to the paternal allele is indicative of microsatellite instability. As previously mentioned, microsatellite instability can be indicative of a mutation at the locus in which the microsatellite resides. If the microsatellite region is associated with a tumor suppressor gene or an oncogene, the detection of microsatellite instability in an allele in a subpopulation of cells in a biological sample is indicative of the potential for cancer or that cancer or precancer may have already developed. Further testing as described herein (either by invasive or noninvasive means) may then be conducted.

In an alternative embodiment, a "fingerprint" of microsatellites is taken from regions associated with cancer-causing genes in a sample obtained from a patient. Such a fingerprint may be obtained by standard methods. The fingerprint comprises the sequence of wild-type microsatellites associated with the cancer-causing gene or genes. Once obtained, the fingerprint is stored and is used in future tests of samples from the same patient in order to monitor changes in microsatellite regions (i.e. microsatellite instability) that may be associated with the development of cancer. Changes in microsatellite length and/or sequence over time may be used to prescribe additional testing and/or treatment in order to detect and remove cancerous tissue at an early stage in its etiology.

Additional embodiments of the invention are apparent upon consideration of the following claims.

- 28 -

CLAIMS

What is claimed is:

- 1 1. A method of screening for the presence of a disease in a patient, the method
2 comprising the steps of:
 - 3 a) obtaining a sample comprising a circumferential surface of a stool
4 voided by a patient; and
 - 5 b) performing an assay to detect in the sample characteristics
6 indicative of the presence of said disease.
- 1 2. A method for screening for the presence of a colorectal cancer or precancerous
2 lesion in a patient, the method comprising the steps of:
 - 3 a) obtaining a stool voided by a patient;
 - 4 b) removing a circumferential surface from said stool; and
 - 5 c) performing an assay on said surface to detect characteristics
6 indicative of the presence of cells or cellular debris shed onto said
7 stool.
- 1 3. The method of claim 2, wherein said assay detects debris from a clonal
2 population of transformed cells comprising said lesion.
- 1 4. The method of claim 3, wherein said assay detects a protein expressed by the
2 transformed cells indicative of the presence of said cells.
- 1 5. The method of claim 3, wherein said assay detects a DNA characteristic
2 indicative of the presence of said cells.
- 1 6. The method of claim 1, further comprising the step of homogenizing said portion
2 in a physiologically compatible buffer prior to step (b).
- 1 7. The method of claim 6, wherein said physiologically compatible buffer comprises
2 a detergent and a proteinase.
- 1 8. The method of claim 1, wherein said assay detects the presence of
2 carcinoembryonic antigen shed from said lesion.

- 29 -

- 1 9. The method of claim 1, wherein said assay comprises the step of exposing said
2 sample to an antibody which specifically binds a molecule characteristic of the
3 presence of said debris.
- 1 10. The method of claim 5, wherein said DNA characteristic is a mutation.
- 1 11. The method of claim 10, wherein said mutation is selected from the group
2 consisting of loss of heterozygosity and microsatellite instability.
- 1 12. The method of claim 10, wherein said characteristics comprise a deletion in a
2 tumor suppressor allele.
- 1 13. The method of claim 1, wherein said assay comprises the step of determining
2 whether a difference exists in said sample between a number X of a first allele
3 known or suspected to be mutated in a subpopulation of cells in the sample and
4 a number Y of a second allele that is known or suspected not to be mutated in a
5 subpopulation of cells in the sample, the presence of a statistically-significant
6 difference being indicative of a mutation in a subpopulation of cells in the
7 sample and the potential presence of a cancerous or precancerous lesion.
- 1 14. The method of claim 1, wherein said assay comprises the step of determining
2 whether a difference exists between a number of a target tumor suppressor
3 allele in the sample and a number of a non-cancer-associated reference allele in
4 the sample, the presence of a statistically-significant difference being indicative
5 of a deletion of the target tumor suppressor allele in a subpopulation of cells in
6 the sample and the potential presence of a cancerous or precancerous lesion.
- 1 15. The method of claim 5, wherein said characteristics comprise a loss of
2 heterozygosity encompassing a polymorphic locus.
- 1 16. The method of claim 13, wherein said assay comprises the steps of:
2 a) detecting an amount of a maternal allele at a polymorphic locus in
3 the biological sample;
4 b) detecting an amount of a paternal allele at the polymorphic locus in
5 the biological sample; and

- 30 -

- 6 c) determining whether a difference exists between the amounts of
7 maternal and paternal allele,
8 the presence of a statistically-significant difference being indicative of a deletion
9 at the polymorphic locus in a subpopulation of cells in the biological sample and
10 the potential presence of a lesion.
- 1 17. The method of claim 1, comprising the additional step of performing a visual
2 examination of the colon of a patient presenting positive assay results.
- 1 18. The method of claim 1, wherein said at least cross-sectional portion comprises
2 an entire stool voided by a patient.
- 1 19. A method for reducing morbidity from colorectal cancer in a population, the
2 method comprising the steps of:
- 3 a) obtaining a sample comprising at least a representative portion of
4 a stool voided by a patient;
- 5 b) performing an assay to detect in the sample the presence of
6 cellular debris shed from the lesion into the voided stool; and
7 c) performing a visual examination of the colon of a patient presenting
8 positive results in said assay to detect the presence of a lesion.
- 1 20. The method of claim 19, wherein said assay detects a loss of DNA in a portion of
2 debris shed from a clonal population of transformed epithelial cells comprising
3 said lesion.




FIG. 1




FIG. 2




FIG. 3

SUBSTITUTE SHEET (RULE26)

4/4

1: 3'ACGCTACGG5'
2: 5'....ATCGGCTTACTGCGATGCC....3'

M

3: 3'....TAGCCGAATGACGCTACGG....5'
4: 5'ATCGGCTTA3'

1: 3'ACGCTACGG5'
2: 5'....ATCGGCTTATTGCGATGCC....3'

F

3: 3'....TAGCCGAATAACGCTACGG....5'
4: 5'ATCGGCTTA3'

FIG. 4

SUBSTITUTE SHEET (RULE26)

SEQUENCE LISTING

<110> Lapidus, Stanley N
Shuber, Anthony P
EXACT LABORATORIES

<120> Methods for Disease Diagnosis From Stool Samples

<130> EXT-015PC

<140>
<141>

<150> US 09/059,713
<151> 1998-04-13

<160> 8

<170> PatentIn Ver. 2.0

<210> 1
<211> 9
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
M1

<400> 1
ggcatcgca 9

<210> 2
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
M2

<400> 2
atcggttac tgcatgcc 19

<210> 3
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
M3

<400> 3
ggcatcgcag taagccgat 19

<210> 4
<211> 9
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
M4

<400> 4
atcggtta 9

<210> 5
<211> 9
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
F1

<400> 5
ggcatcgca

9

<210> 6
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
F2

<400> 6
atcggcttat tgcgatgcc

19

<210> 7
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
F3

<400> 7
ggcatcgcaa taagccgat

19

<210> 8
<211> 9
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
F4

<400> 8
atcggctta

9

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/08013

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 G01N33/50 C12Q1/68 G01N33/72 G01N33/574

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 G01N C12Q B01L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, L	WO 97 28450 A (EXACT LABORATORIES) 7 August 1997 (1997-08-07) the whole document	1-20
X	EP 0 391 565 A (HELENA LABORATORIES CORPORATION) 10 October 1990 (1990-10-10) column 8, line 15 - column 29; claims	1, 18
A	US 4 578 358 A (N.H.OKSMAN) 25 March 1986 (1986-03-25) abstract; claims column 7, line 5 - line 11	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 September 1999

Date of mailing of the international search report

23/09/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Luzzatto, E

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/08013

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9728450 A	07-08-1997	US 5741650 A	21-04-1998
		AU 704696 B	29-04-1999
		AU 2240697 A	22-08-1997
		CA 2215263 A	07-08-1997
		EP 0817968 A	14-01-1998
EP 391565 A	10-10-1990	US 5192501 A	09-03-1993
		CA 2012830 A	04-10-1990
		DE 69026859 D	13-06-1996
		DE 69026859 T	12-09-1996
		JP 2295495 A	06-12-1990
		US 5196167 A	23-03-1993
US 4578358 A	25-03-1986	US 5217874 A	08-06-1993
		AU 547546 B	24-10-1985
		AU 2470984 A	08-11-1984
		BR 8400933 A	05-03-1985
		CA 1212891 A	21-10-1986
		EP 0124215 A	07-11-1984
		JP 1688475 C	11-08-1992
		JP 3054305 B	19-08-1991
		JP 59204766 A	20-11-1984
		ZA 8401079 A	26-09-1984

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 33/50, C12Q 1/68, G01N 33/72, 33/574	A1	(11) International Publication Number: WO 99/53316 (43) International Publication Date: 21 October 1999 (21.10.99)
(21) International Application Number: PCT/US99/08013 (22) International Filing Date: 13 April 1999 (13.04.99) (30) Priority Data: 09/059,713 13 April 1998 (13.04.98) US (71) Applicant: EXACT LABORATORIES, INC. [US/US]; 63 Great Road, Maynard, MA 01754 (US). (72) Inventors: LAPIDUS, Stanley, N.; 12 Old Evergreen Road, Bedford, NH 03110 (US). SHUBER, Anthony, P.; 11 Grant Street, Milford, MA 01757 (US). (74) Agent: WALLER, Patrick, R., H.; Testa, Hurwitz & Thibault, LLP, High Street Tower, 125 High Street, Boston, MA 02110 (US).	(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: METHODS FOR DISEASE DIAGNOSIS FROM STOOL SAMPLES (57) Abstract The present invention provides methods for preparing a stool sample in order to screen for the presence of indicators of a disease, for example a subpopulation of cancerous or precancerous cells. The methods take advantage of the recognition that cellular debris from cancerous and precancerous cells is deposited onto only a longitudinal stripe of stool as the stool is forming in the colon. Accordingly, methods of the invention comprise obtaining a representative sample, such as a circumferential or cross-sectional sample of stool in order to ensure that any disease indicator, such as cellular debris that is shed by colonic cells, is obtained in the sample.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHODS FOR DISEASE DIAGNOSIS FROM STOOL SAMPLES

This patent application is a continuation-in-part of U.S. Serial Number 08/699,678, filed August 14, 1996, the disclosure of which is incorporated herein by reference.

FIELD OF THE INVENTION

5 This invention relates to methods for disease diagnosis, including the early detection of colon cancer in patients, and more particularly to methods for preparing stool samples for disease diagnosis, including the detection of colon cancer, so as to assure or increase the likelihood that the sample will contain the diagnostically relevant information if the patient has a disease, for example a cancerous or precancerous
10 lesion, and to methods for stool sample analysis.

BACKGROUND OF THE INVENTION

 Stool samples frequently must be prepared for medical diagnostic analysis. Stool samples may be analyzed to help diagnose medical conditions ranging from parasitic, bacterial or viral infections to inflammatory bowel disease and colorectal
15 cancer.

 Colorectal cancer is a leading cause of death in Western society. However, if diagnosed early, it may be treated effectively by surgical removal of the cancerous tissue. Colorectal cancers originate in the colorectal epithelium and typically are not extensively vascularized (and therefore not invasive) during the early stages of
20 development. Colorectal cancer is thought to result from the clonal expansion of a single mutant cell in the epithelial lining of the colon or rectum. The transition to a highly vascularized, invasive and ultimately metastatic cancer which spreads throughout the body commonly takes ten years or longer. If the cancer is detected prior to invasion, surgical removal of the cancerous tissue is an effective cure.
25 However, colorectal cancer is often detected only upon manifestation of clinical symptoms, such as pain and black tarry stool. Generally, such symptoms are present only when the disease is well established, often after metastasis has occurred, and the

- 2 -

prognosis for the patient is poor, even after surgical resection of the cancerous tissue. Early detection of colorectal cancer therefore is important in that detection may significantly reduce its morbidity.

Invasive diagnostic methods such as endoscopic examination allow for direct
5 visual identification, removal, and biopsy of potentially cancerous growths such as polyps. Endoscopy is expensive, uncomfortable, inherently risky, and therefore not a practical tool for screening populations to identify those with colorectal cancer. Non-invasive analysis of stool samples for characteristics indicative of the presence of colorectal cancer or precancer is a preferred alternative for early diagnosis, but no
10 known diagnostic method is available which reliably achieves this goal.

Current non-invasive diagnostic methods involve assaying stool samples for the presence of fecal occult blood or for elevated levels of carcinoembryonic antigen, both of which are suggestive of the presence of colorectal cancer. Additionally, recent developments in molecular biology provide methods of great potential for detecting the
15 presence of a range of DNA mutations or alterations associated with and indicative of the presence of colorectal cancer. The presence of such mutations theoretically can be detected in DNA found in stool samples during the early stages of colorectal cancer. However, stool comprises cells and cellular debris from the patient, from microorganisms, and from food, resulting in a heterogeneous population of cells. This
20 makes detection of a small, specific subpopulation impossible to detect reliably.

Stool diagnostic assays for colorectal cancer described in the art typically are performed on samples prepared from randomly sampled portions of voided stool. However, samples prepared according to such methods do not reproducibly yield characteristics indicative of the presence of colorectal cancer or precancer, even when
25 prepared from stool voided by a patient with colorectal cancer or precancer. There is therefore a need in the art for methods for early diagnosis of colorectal cancer or precancer that will reproducibly detect characteristics indicative of the presence of cancerous or precancerous material in samples prepared from stool voided by a patient with colorectal cancer or precancer. Such methods are provided herein.

- 3 -

SUMMARY OF THE INVENTION

It has now been appreciated that cells and cellular debris are shed from colonic epithelial cells onto forming stool in a longitudinal "stripe" of material along the length of the stool. The shed material is confined to this longitudinal stripe, as shown in

5 Figure 1 (designated "C"). Based upon this recognition, Applicants teach that stool sample preparation for diagnostic testing must include taking a representative sample in order to ensure that the sample will contain any cells or cellular debris that was shed into the stool as it passed through the colon. Accordingly, methods of the invention

10 comprise obtaining at least a representative (e.g. a cross-section or circumferential surface) portion of stool voided by a patient, and performing an assay to detect in the sample the presence of cells or cellular debris shed from epithelial cells lining the colon that may be indicative of cancer or precancer. Most often, such cells will be derived from a polyp or a cancerous or precancerous lesion at a discrete location along the colon. For purposes of the present invention, a precancerous lesion comprises

15 precancerous cells, and precancerous cells are cells that have a mutation that is associated with cancer and which renders such cells susceptible to becoming cancerous. As shown in Figure 1, a cross-sectional sample is a sample that contains at least a circumferential surface of the stool (or portion of a stool comprising an entire cross-sectional portion), as, for example, in a coronal section or a sagittal section. A

20 sample comprising the surface layer of a stool (or of a cross-section of a stool) also contains at least a circumferential surface of the stool. Both cross-sections and circumferential surfaces comprise longitudinal stripes of sloughed colonic epithelium, and are therefore representative samples.

In a preferred embodiment, methods of the invention comprise the steps of

25 obtaining at least a circumferential surface or cross-sectional portion of a stool voided by a patient, and performing an assay to detect debris indicative of disease. For example, such debris may comprise a clonal subpopulation of cells having one or more mutations (for purposes of the present application, a mutation is a deletion, substitution, addition, modification, intercalation or rearrangement of DNA). Preferred

30 methods of the invention comprise detection of characteristics of such transformed cells, including, for example, mutations, proteins expressed uniquely or in altered

- 4 -

amounts in transformed cells, and blood. Particularly preferred methods of the invention comprise obtaining at least a circumferential surface or cross-sectional portion of a stool sample, and performing an assay to detect DNA characteristics indicative of the presence of a clonal subpopulation of cells in the sample. The clonal
5 subpopulation may be, for example, a subpopulation of cancerous or precancerous cells, having a mutation in, for example, a p53 tumor suppressor gene. Clonal subpopulations of cells detected by methods according to the invention are often characterized by a massive loss of DNA, resulting in a loss of heterozygosity that renders ineffective the gene or genes encompassed by the deletion. Alternative
10 methods of the invention comprise performing an assay to detect the presence of an infectious disease, for example debris from a microorganism.

Methods of the invention also comprise obtaining a representative (*i.e.*, cross-sectional or circumferential) sample of stool and homogenizing the stool in a buffer, such as a buffer comprising a detergent and a proteinase and optionally a DNase
15 inhibitor.

In methods according to the invention, an assay performed on at least a circumferential surface or cross-sectional portion of stool may be an assay to detect a disease when a diagnostic indicator of the disease is incorporated into stool. Methods of the invention are useful to detect cellular debris shed from the epithelial lining of the colon, and from other tissue sources that shed cellular material into the gastrointestinal tract and preferably into the colon. Shed material may be from colonic epithelial cells, pancreatic cells, or may be indicative of infection (e.g. bacterial or viral nucleic acids or proteins). In methods according to the invention, an assay performed on at least a circumferential surface or cross-sectional portion of stool may be an assay to detect the
20 presence of elevated levels of carcinoembryonic antigen shed from cells lining the colon. Such an assay may also comprise detecting the presence of occult blood. However, methods of the invention preferably comprise an assay wherein the sample is exposed to an antibody that specifically binds to a molecule characteristic of cellular debris shed from cells comprising a subpopulation of cells having a mutation that is
25 potentially associated with cancer.
30

- 5 -

Methods of the invention are especially and most preferably useful for detecting DNA characteristics indicative of a subpopulation of transformed cells in a representative stool sample. The DNA characteristics may be, for example, mutations, including loss of heterozygosity, microsatellite instability, and others. An assay for DNA characteristics in a method of the invention may comprise the step of determining whether a difference exists in a number X of a first allele known or suspected to be mutated in a subpopulation of cells in a representative stool sample, and a number Y of an allele known or suspected not to be mutated in the sample, a statistically-significant difference being indicative of a mutation and the possible presence of cancer in a subpopulation of cells in the sample. In an embodiment of the invention, the difference between a number of a tumor suppressor gene and a number of a non-cancer-associated gene are compared, a statistically-significant difference in the numbers being indicative of a mutation in the tumor suppressor gene.

Assays useful in the practice of methods according to the invention also include an assay to detect the presence of a deletion or other mutation in a region encompassing a polymorphic nucleotide. In such an assay, a number of a polymorphic nucleotide present at maternal and paternal alleles, wherein the patient is heterozygous for the polymorphic nucleotide, is determined. A statistically significant difference between a number of a polymorphic nucleotide in a maternal allele and a paternal allele is indicative of the presence of a deletion in one of the two alleles. Another useful assay is one to detect an infectious disease.

Methods of the invention typically include, following sample preparation and an assay for characteristics of cells or cellular debris, a visual examination of the colon in order to determine if a polyp or other lesion is, in fact, present. Finally, surgical resection of abnormal tissue may be done in order to prevent the spread of cancerous or precancerous tissue.

Accordingly, methods of the invention provide means for screening for the presence of a cancerous or precancerous subpopulation of cells in a heterogeneous sample, such as a stool sample. Methods of the invention reduce morbidity and mortality associated with lesions of the colonic epithelium. Moreover, methods of the invention comprise more accurate screening methods than are currently available in

- 6 -

the art, because current methods take advantage of the observation that cancerous or precancerous cells shed debris only onto or into part of the surface of the forming stool. The present methods reliably assay over the entire circumference of the stool, thereby increasing the likelihood of detecting an abnormality if one exists. Further aspects and advantages of the invention are contained in the following detailed description thereof.

DESCRIPTION OF THE DRAWINGS

Figure 1 is a diagram of a cylinder which represents a formed stool and shows various cross-sections which will contain material from the entire circumference of a stool. The Section labeled "A" is a typical coronal section and the section labeled "B" is a typical sagittal section. The strip labeled "C" represents material shed from cancerous tissue which is deposited in a longitudinal stripe.

Figure 2 is a schematic diagram of a receptacle for containing a stool sample.

Figure 3 is a schematic diagram of a multi-orifice impedance counter; wherein reference numeral 1 indicates the direction of flow through the column; reference numeral 2 indicates a plunger means for forcing material downward in the column; reference numerals 3 and 4 are different-sized hybridization beads; reference numeral 5 is an optional filter for extracting unwanted particles; reference numeral 6 indicates an array of orifices for measuring differential impedance; and reference numeral 7 is a collection chamber.

Figure 4 is a diagram showing primers useful for the detection of single base polymorphisms.

DETAILED DESCRIPTION OF THE INVENTION

Methods according to the present invention are useful for the preparation of stool samples that will reproducibly contain cells or cellular debris shed from a clonal population of cancerous or precancerous cells, if such a population is present at any site along the colon of a patient. These samples are then used to perform assays to detect characteristics indicative of cancer in a highly-reproducible and accurate way. In preferred methods of the invention, a sample representative of the surface of a stool is obtained. Such methods provide an improvement over the art inasmuch as they teach removing the portion of stool onto which the colonic epithelium sheds debris.

- 7 -

Two preferred methods for obtaining such a portion are cross-sectioning and extracting the surface of the stool. In a more preferred method, the surface of a cross-section is obtained. Without the recognition that at least a cross-sectional sample must be obtained, there is no means for reproducibly obtaining a sample that will contain a
5 cancerous or precancerous subpopulation of cells, if one exists.

Methods described in the art do not recognize that, unlike infection by parasites, bacteria and viruses, characteristics indicative of the presence of colon cancer, especially early stage colon cancer, are found only in a specific portion of voided stool. If the sampled portion of stool does not include the portion that happens to contain
10 cells and cellular debris shed from early-stage cancer tissue, the diagnostic assay necessarily will fail to detect the characteristics indicative of the presence of colorectal cancer in a reliable manner even if homogenized, i.e., will produce a false-negative result. Methods of the invention are also useful to detect the presence of bacteria, virus or parasites, as they will also be present on the surface of formed stool.

15 Sloughed cells from, for example, a polyp forming in the epithelial lining of the colon, or on early stage cancerous lesions, are sloughed onto only the portion of the forming stool that comes into contact with the polyp or lesion. Accordingly, in early stage disease, only a small portion of the surface layer of the forming stool will contain sloughed cells, and if that portion happens not to be taken as part of the sample, an
20 assay for indicia of colon cancer necessarily will produce a false-negative result. A brief review of the anatomy and physiology of the colon will aid in an understanding of this phenomenon.

A typical adult colon is approximately six feet in length, with a diameter of about two to three inches. Numerous bends and folds are present throughout its length. The
25 colon removes water from liquid or semi-liquid waste material that enters the colon, and relatively solid stool begins to form in the proximal third of the colon. Epithelial cells line the lumen of the colon, and the luminal surface is organized into microscopic crypts. Colorectal epithelial cells are replaced every four to five days. The epithelial cells divide rapidly at the base of the crypts and migrate to the apices, where cells
30 appear to undergo apoptosis (programmed cell death), and cellular debris is shed into the lumen. The lining of the colorectal lumen is elastic and the diameter of the lumen is

- 8 -

determined by the volume of stool that is passing through the colon at any given time. As a result, the surface of the forming stool passing through the colon is in direct contact with the epithelial lining of the lumen. Shed epithelial cells (which may or may not have undergone apoptosis) and cellular debris therefore are incorporated onto the surface of stool as it passes through the colon.

Cells and cellular debris from colorectal epithelial cancers therefore also are shed onto forming stool. Most colorectal cancers develop in regions of the colon where stool is relatively solid, indeed approximately one third of such cancers develop in the rectum. Markers indicative of the presence of cancer, including cells, cellular debris, DNA, blood, and carcinoembryonic antigen, are shed onto the portion of the forming stool that contacts the cancerous tissue as the stool passes through the colon. Since the stool is relatively solid, these markers will remain on or near the surface of the stool where they were deposited and will not be homogeneously dispersed throughout the stool. As stool passes over a cancerous or precancerous growth, material from the growth will be deposited along the stool, but only on the part of the stool circumference that comes into direct contact with the cancerous or precancerous tissue comprising the lesion. Stool voided by a patient with colorectal cancer or precancer is therefore characterized by a longitudinal "stripe" of diagnostically relevant material derived from the cancerous or precancerous tissue.

A sample that does not include material from the entire circumference of a stool voided by a patient with colorectal cancer or precancer will not reproducibly contain material derived from the cancerous or precancerous tissue. Currently, random, non-cross-sectional samples ("smears") of voided stool are analyzed in clinical settings. In these, sloughed cancerous or precancerous cells and cellular debris have no possibility of detection unless the sample happens by chance to contain the portion of stool which made contact with the region of the colon from which cells were sloughed.

Furthermore, cancers typically develop by clonal expansion of a single mutant cell, and in the early stages of the disease, i.e., when surgical removal is an effective cure, the cancerous lesion will be very small and may lie on a small arc of the circumference of the colon. Material derived from such an early stage cancer therefore will be shed onto or into stool in a very narrow stripe (labeled C in Figure 1).

- 9 -

Consequently, a sample that does not contain the entire circumference of a stool voided by a patient with early stage colorectal cancer or precancer only by chance will contain material indicative of the presence of the early stage cancerous or precancerous condition. However, early detection of colorectal cancer is very
5 important for effective surgical intervention. The present invention provides methods for reproducible early detection of characteristics indicative of the presence of cancer or precancer in a patient.

Similarly, the present invention provides methods for reproducibly detecting an indicator of the presence of an infectious agent, such as a virus, bacterium, parasite or
10 other microorganism, in stool from an infected patient. Methods of the invention are particularly useful for reproducibly detecting an infectious agent, or an indicator characteristic thereof, that is deposited onto narrow area of stool as it forms in the colon.

Analysis of at least a circumferential surface or cross-sectional sample of stool
15 ensures that a disease indicator, incorporated on a stool as it moves through the colon, will be present in a diagnostic assay sample. For example, analysis of at least a circumferential surface or cross-sectional sample of stool (including a whole stool), as shown in Figure 1, ensures that at least a portion of cells and cellular debris shed from any existing cancerous or pre-cancerous cells (even if shed from small early stage
20 cancerous or pre-cancerous tissue, e.g., small polyps) will be present in the portion of the stool sample to be analyzed. Indeed taking at least a circumferential surface or cross-section of the stool sample avoids the possibility of analyzing stool portions that will not contain sloughed cancerous or precancerous cells even when the patient has colorectal cancer or precancer.

25 Once a circumferential surface or cross-sectional sample of stool is obtained, it may be homogenized by known methods to distribute cells and cellular debris throughout the sample. An assay then is performed on the homogenate, or an extract of the homogenate, to detect the presence of cells and/or cellular debris in the sample. The assay may be any one or a combination of histological cellular assays, antibody
30 based immunoassays (or other formats) designed to detect the presence of a molecule characteristic of transformation such as a protein, or DNA-based assays for detecting

- 10 -

mutations or genetic characteristics indicative of colorectal cancer. Known assay protocols, those disclosed herein or in U.S. Patent No. 5,670,325, or assays hereafter developed may be used in the practice of the invention. Non-limiting examples of useful known assay protocols include those disclosed in U.S. Patent Nos. 5,137,806
5 (detection of sequences in selected DNA molecules), 5,348,855 (assay for nucleic acid sequences), 5,512,441 (detection of mutant alleles), 5,272,057 and 5,380,645 (RFLP analysis), 5,527,676 (detection of p53 gene sequences), 5,330,892 (detection of MCC gene sequences), 5,352,775 (detection of APC gene sequences), 5,532,108 (detection of DCC gene sequences), and in WO96/08514 (monoclonal antibodies against human
10 colon carcinoma-associated antigens), the disclosures of each of which are incorporated by reference herein. Alternatively, or additionally, an assay for fecal occult blood may be performed as reported in U.S. Patent Nos. 4,333,734 and 5,196,167, incorporated by reference herein. Assays useful in the context of the present invention also include an assay for carcinoembryonic antigen as reported in
15 U.S. Patent No. 5,380,647, incorporated by reference herein. In addition, assays described in USSN 08/808,763; 08/815,576; 08/876,857; 08/876,638; 08/877,333; and 60/063,219; incorporated by reference herein, are also useful in the context of the present invention. Finally, the sample may be prepared, as reported in U.S. Patent No. 4,857,300, incorporated by reference herein, for histological examination to detect
20 characteristics indicative of the presence of cancerous or precancerous cells.

The purpose of any assay protocol used in connection with obtaining at least a cross-sectional sample is to identify candidates for subsequent invasive diagnostic procedure such as colonoscopy or sigmoidoscopy. The assay accordingly need not definitively detect the presence of a cancerous or precancerous lesion, although false
25 negatives obviously are to be avoided. The goal of the test protocol is not to determine whether in the vast quantities of cell debris in the sample there are a few cells which bear a mutation commonly associated with early stage transformation, but rather whether the sample contains debris indicative of clonal expansion of a mutant cellular subpopulation. Maximum benefit will come from an assay designed to detect the likely
30 presence of a clonally expanded cell population, i.e., of transformed colonic epithelial cells which comprise the cancerous or precancerous lesions. Assays having the ability

- 11 -

to detect cancer or precancer in early stages are preferred in methods of the invention. Assays using polymerase chain reaction (PCR), restriction fragment length polymorphism (RFLP) or other methods for nucleic acid analysis may be used to detect known DNA characteristics indicative of the presence of colorectal cancer or
5 precancer. More precise methods for quantitative detection of cellular debris, such as DNA fragments or segments, may be used to analyze cross-sectional samples according to methods described herein.

A preferred assay interrogates the sample for DNA characteristics indicative of the development of cancer or precancer. However, assays for use with methods of the
10 invention may detect any abnormal cellular debris shed from clinically-relevant transformed tissue. Thus, in accordance with a preferred aspect of the invention, an assay is used to detect the presence of characteristics of cells which have experienced loss of heterozygosity, microsatellite instability or other mutation.

The following examples provide details of methods according to the invention.
15 However, numerous additional aspects of the invention, especially in terms of assays to be performed, will become apparent upon consideration of the following detailed description thereof.

Example 1

Preparation of a Stool Sample

20 A sample is prepared such that it contains at least a circumferential surface or cross-sectional portion of a stool voided by a patient.

Stool is voided into a receptacle that is preferably small enough to be transported to a testing facility. The receptacle may be fitted to a conventional toilet such that the receptacle accepts stool voided in a conventional manner. The
25 receptacle may comprise a mesh or screen of sufficient size and placement such that stool is retained while urine is allowed to pass through the mesh or screen and into the toilet. The receptacle additionally may comprise means for removing a circumferential surface or cross-sectional portion from the stool. Moreover, the receptacle may comprise means for introducing homogenization buffer or one or more preservatives,
30 such as alcohol, a solution of high salt concentration, antibiotics, and chaotropic salts in order to neutralize bacteria present in the stool sample. The homogenization buffer

- 12 -

may be a physiologically compatible buffer such as phosphate buffered saline, and may comprise salt such as 20-100 mM NaCl or KCl. The homogenization buffer may also comprise a detergent, such as 1-10% SDS or triton, and/or a proteinase, such as proteinase K. The buffer may also contain inhibitors of DNA and RNA degrading enzymes.

The receptacle, whether adapted to fit a toilet or simply adapted for receiving the voided stool sample, should include sealing means sufficient to contain the voided stool sample and any solution added thereto and to prevent the emanation of odors. An exemplary receptacle is shown in Figure 2. As shown in that figure, the receptacle has a support frame 1 which is placed directly over the toilet bowl 2. The support frame 1 has attached thereto an articulating cover 3 which may be placed in a raised position, as shown in figure 2, for depositing of sample or a closed position (not shown) for sealing voided stool within the receptacle. The support frame 1 additionally has a central opening 4 traversing from a top surface 5 through to a bottom surface 6 of the support frame 1. The bottom surface 6 directly communicates with a top surface 7 of the toilet 2. Extending from the bottom surface 6 of the support frame 1 is a means 8 for capturing voided stool. Means 8 may be fixedly attached to the support frame 1 or may be removably attached for removal subsequent to deposition of stool. Means 8 may comprise a further means for removing at least a circumferential surface or cross-sectional portion from the voided stool. A preferable sample size is at least 5-10 g or at least 5-10 ml. A means to assess the presence of a minimal sample size may comprise a physical diagram indicating the minimal sample size. Alternatively a means to assess the presence of a minimal sample size may comprise the displacement of a liquid or of a mechanical device to a minimal level upon deposit of the stool sample.

In a preferred embodiment, a cross-sectional portion is removed from the voided stool by making one or more sagittal or coronal section through the stool, as shown in Figure 1. The removed portion comprises material from an entire circumferential surface of the stool. Alternatively, a circumferential surface of a stool is prepared by taking the entire surface layer of the stool, or the entire surface layer of a cross-sectional portion of the stool. In a preferred embodiment, the surface layer is approximately 0.001 to 0.1 mm thick. In another preferred embodiment, the surface

- 13 -

layer is approximately 0.1 to 20 mm thick. In a more preferred embodiment, the surface layer is approximately 5 mm thick. In a most preferred embodiment, the surface layer is representative of material deposited on the forming stool, and reproducibly contains markers or indicators, such as shed cellular material, of a disease that is present in the

5 patient. In one embodiment, a stool surface layer is obtained by cutting or scraping off the surface of the stool or cross-sectional portion of the stool. In a preferred embodiment, a stool surface layer is prepared using a cylindrical cutting device (for example a rigid loop) that has a diameter that is smaller than the diameter of the stool or cross-sectional portion of the stool. The surface layer is obtained when the stool

10 sample is forced through the cutting device. In a preferred embodiment, the diameter of the cutting device is adjustable. Alternatively, a circumferential surface of a stool is obtained by contacting the surface of a stool or cross-sectional portion of a stool with a sampling device. In one embodiment, the sampling device is first wrapped around the stool sample and subsequently removed or peeled off. The layer of stool that is peeled

15 off with the sampling device is then processed as described herein. Alternatively, the stool sample is rolled along the surface of the sampling device, and the layer of stool that adheres to the surface is then processed as described herein. Alternatively, a diagnostic assay is performed directly on the layer of material on the sampling device. It is important that the entire circumference of the stool is sampled. In a preferred

20 embodiment, the sampling device has a surface that stool adheres to. In a more preferred embodiment, the sampling device surface preferentially binds diagnostic markers or indicators of a disease (for example nucleic acid or protein molecules), and thereby removes them from the surface of the stool sample. In one embodiment, the sampling device is an adherent paper or membrane. In a preferred embodiment, a

25 stool is frozen prior to processing. In one embodiment the stool is frozen to between -5 and -30 °C. In a preferred embodiment the cross-sectional portion or circumference of the stool is obtained from the frozen or partially thawed stool. In an alternative embodiment, a whole stool may be used for diagnostic assays. The stool or portion of stool that is obtained contains sufficient material to allow subsequent diagnostic assays

30 to be performed.

- 14 -

Once obtained, the entire circumference or cross-sectional sample of stool is homogenized in an appropriate buffer, such as phosphate buffered saline. Homogenization means and materials for homogenization are generally known in the art. Thus, particular homogenization methods may be selected by the skilled artisan and may depend upon the assay to be employed. The buffer may contain detergent, salt, proteinase, inhibitors of DNA and RNA degrading enzymes. The composition of the buffer will depend on the type of assay to be performed. If a fecal occult blood assay is to be performed, the buffer may contain chemical compounds which react with blood to produce a color, the intensity of which can be measured. Buffers useful for detecting the presence of fecal occult blood are known in the art. If a test is to be performed for the presence of a particular protein, the buffer should not contain a proteinase capable of degrading such tumor marking antigens.

DNA or RNA may be isolated from the homogenate using methods known in the art. Subsequent tests may be performed on the isolated DNA and RNA.

Example 2

Exemplary Enumerative methods for Detection of Colorectal Cancer or Precancer in stool samples

DNA characteristics associated with the presence of colorectal cancer or a precancerous lesion may be detected in stool samples prepared according to the invention, using, for example, the methods described in the following sections. A careful endoscopic examination preferably is performed on positive Individuals, followed by early surgical excision of any diseased tissue.

A. Reference-Target

Methods of the invention are used to prepare a stool sample followed by detection of a deletion or other mutation in the p53 tumor suppressor gene. The p53 gene is a good choice because a loss of heterozygosity in p53 is often associated with colorectal cancer. An mRNA sequence corresponding to the DNA coding region for p53 is reported as GenBank Accession No. M92424. At least a cross-section of a voided stool sample is obtained and prepared according to methods of the invention as described immediately above. The sample need not be further processed for analysis. However, DNA or RNA may optionally be isolated from the sample according to

- 15 -

methods known in the art. See, Smith-Ravin, *et al.*, *Gut*, 36: 81-86 (1995), incorporated by reference herein.

Nucleic acids may be sheared or cut into small fragments by, for example, restriction digestion. The size of nucleic acid fragments produced is not critical, subject
5 to the limitations described below. A target allele that is suspected of being mutated (p53 in this example) and a reference allele are chosen. A reference allele may be any allele known normally not to be mutated in colon cancer.

Either portions of a coding strand or its complement may be detected. For exemplification, detection of the coding strand of p53 and reference allele are
10 described herein. Complement to both p53 and reference allele are removed by hybridization to anti-complement oligonucleotide probes (isolation probes) and subsequent removal of duplex formed thereby. Methods for removal of complement strands from a mixture of single-stranded oligonucleotides are known and include techniques such as affinity chromatography. Upon converting double-stranded DNA to
15 single-stranded DNA [See, e.g., Sambrook, *et al.*, *Molecular Cloning, A Laboratory Manual* (1989) incorporated by reference herein], sample is passed through an affinity column packed with bound isolation probe that is complementary to the sequence to be isolated away from the sample. Conventional column chromatography is appropriate for isolation of complement. An affinity column packed with sepharose or other
20 appropriate materials with attached complementary nucleotides may be used to isolate complement DNA in the column, while allowing DNA to be analyzed to pass through the column. See Sambrook, *Supra*. As an alternative, isolation beads may be used to exclude complement as discussed in detail below.

After removal of complement strands, first oligonucleotide probes which
25 hybridize to at least a portion of the p53 allele and second oligonucleotide probes that hybridize to at least a portion of the reference allele are obtained. The probes are labeled with a detectable label, such as fluorescein or with detectable particles. Distinct labels for the probes are preferred. However, identical labels may be used if, for example, sample is assayed in two separate aliquots. Probes may be labeled with
30 identical or with distinct labels. However, distinct labels are preferred.

- 16 -

Labeled probes then are exposed to sample under hybridization conditions. Such conditions are well-known in the art. See, e.g., Wallace, et al., *Nucleic Acids Res.*, 6:3543-3557 (1979), incorporated by reference herein. First and Second oligonucleotide probes that are distinctly labeled (i.e. with different radioactive isotopes, fluorescent means, or with beads of different size, See *infra*) are applied to a single aliquot of sample. After exposure of the probes to sample under hybridization conditions, sample is washed to remove any unhybridized probe. Thereafter, hybridized probes are detected separately for p53 hybrids and reference allele hybrids. Standards may be used to establish background and to equilibrate results. Also, if differential fluorescent labels are used, the number of probes may be determined by counting differential fluorescent events in a sample that has been diluted sufficiently to enable detection of single fluorescent events in the sample. Duplicate samples may be analyzed in order to confirm the accuracy of results obtained.

If there is a statistically-significant difference between the amount of p53 detected and the amount of the reference allele detected, it may be assumed that a mutation has occurred in p53 so as to alter its sequence and prevent hybridization of the probe, or that at least a portion of the region of the genome containing p53 has been lost in a subpopulation of cells shed from the colon. The patient therefore may be at risk for developing or may have developed colon cancer. Statistical significance may be determined by any known method. See, e.g., Steel, et al., *Principles and Procedures of Statistics: A Biometrical Approach* (McGraw, Hill, 1980). Statistical methods are also outlined in co-owned U.S. Patent No. 5,670,325.

The determination of a p53 mutation allows a clinician to recommend further treatment, such as endoscopy procedures, in order to further diagnose and, if necessary, treat the patient's condition. The following examples illustrate methods that allow direct quantification of hybridization events.

1. Method for Quantitation of Target and Reference Polynucleotides

Enhanced quantification of binding events between hybridization probes and target or reference is accomplished by coupling hybridization probes to particles, such as beads (hybridization beads). In order to obtain a precise quantitative measure of

- 17 -

the amount of a polynucleotide in a sample, hybridization beads are constructed such that each bead has attached thereto a single oligonucleotide probe.

a. Method for Preparation of Probe-Bead Combinations

A single probe is attached to a bead by incubating a large excess of
5 hybridization beads with oligonucleotide probes of a given type (*i.e.*, either first or second oligonucleotide probes). Coupling of probe to bead is accomplished using an affinity-binding pair. For example, beads may be coated with avidin or streptavidin and probes may be labeled with biotin to effect attachment of the probe to the bead. The mixture of beads and probes is agitated such that 100% of the probes are bound to a
10 bead. The mixture is then exposed to a matrix, such as an affinity column or a membrane coated with oligonucleotides that are complementary to the probe. Only beads that have an attached probe will adhere to the matrix, the rest being washed away. Beads with coupled probe are then released from the matrix by melting hybridizations between probe and complement. Multiple exposures to the matrix and
15 pre-washing of the column reduces non-specific binding. Moreover, naked beads (*i.e.*, without attached probe) may be exposed to the matrix to determine a background number of beads that can be expected to attach to the matrix in the absence of probe.

By using a vast excess of beads relative to probe as described above, the vast majority of recovered beads will have only one attached probe. For example, if a
20 mixture has a ratio of 1 probe to 1000 beads, it is expected that only about 1 bead in a million will have two attached probes and even less than one bead in a million will have more than two attached probes. Accordingly, hybridization beads are provided in an effective 1:1 ratio with probe which allows for precise quantification of target and reference polynucleotide as described below.

25 For each assay described below, two distinct hybridization beads are used. A first hybridization bead has attached thereto a single first oligonucleotide probe that is complementary to at least a portion of a target polynucleotide (*e.g.*, a p53 allele). A second hybridization bead, of a size distinct from the first hybridization bead, has attached thereto a single second oligonucleotide probe that is complementary to at
30 least a portion of a reference polynucleotide (*i.e.*, one that is known or suspected not to be mutated in the sample).

- 18 -

b. Use of Beads to Quantify Target and Reference Polynucleotides

5 DNA is melted (denatured to form single-stranded DNA) by well-known methods See, e.g., Gyllenstein, *et al.*, in *Recombinant DNA Methodology II*, 565-578 (Wu, ed., 1995), incorporated by reference herein. One may detect either a coding strand or its complement in order to quantify target and/or reference polynucleotide. For purposes of illustration, the present example assumes detection of the coding strand.

10

2. Removal of Complement

Single-stranded complement of the target polynucleotide (e.g., p53) and reference polynucleotide are removed from the sample by binding to oligonucleotide probes that are complementary to target or reference complement. Such probes, referred to herein as isolation probes, are attached to isolation beads prior to their introduction into the sample. The beads may be magnetized. Thus, when magnetized isolation beads [with attached isolation probe(s)] are introduced into the sample, the attached isolation probes hybridize to complement of target or reference (or vice versa). Isolation beads preferably are introduced in vast excess in order to saturate complement binding. Once hybridization is complete, a magnetic field is applied to the sample to draw the magnetized isolation beads (both with and without hybridized complement) out of the sample. Assuming that a sufficient quantity of isolation beads are introduced into the sample, removal of the isolation beads effectively removes all target and reference complement from the sample.

25 In an alternative method for complement removal, an excess of oligonucleotide probe labeled with biotin is exposed to the melted or dehybridized (single stranded) sample under hybridization conditions. Once hybridization is complete, the sample is exposed to a column containing immobilized avidin. The biotin-labeled probe, whether free or hybridized to complement, is bound by avidin on the column. The remainder of the DNA, including target and reference coding strands to be detected, pass through the column. In contrast to the description of hybridization beads above, beads for

30

- 19 -

removal of complement may each comprise multiple complementary oligonucleotide probes.

3. *Quantitation of Target and Reference*

Two sets of hybridization beads are prepared as described above. Each

5 member of a first set of hybridization beads (all of which are identical to each other) has attached thereto a single oligonucleotide probe that is complementary to at least a portion of the target polynucleotide, i.e., the portion of the genome which is altered in the cells of a cancerous lesion. Each member of a second set of identical hybridization beads (all of which are identical to each other but not to the first set) has attached

10 thereto a single oligonucleotide probe that is complementary to at least a portion of the reference polynucleotide, i.e., a portion of the genome which is not likely to be altered in malignant cells. Members of the second set of hybridization beads are of a size or color distinct from that of members of the first set of hybridization beads. First and second hybridization beads may also be distinguished on the basis of other

15 characteristics. For example, beads may have fluorescent markers that are distinguished by their fluorescence wavelength. Beads with distinct electrochemical charges also may be used. The precise modality used for distinguishing beads is not essential as long as it is possible to distinguish between first and second probe on the basis of distinctions between attached first and second beads.

20 Both sets of hybridization beads are exposed to the sample under hybridization conditions thereby allowing hybridization to reference and target. The sample then is washed to remove unhybridized bead/probe combinations. Unhybridized bead/probe combinations are removed by, for example, passing the sample through a column lined of immobilized DNA complementary to the probe sequence. Thus, any unhybridized

25 bead/probe combinations are retained on the column while duplex passes through. Subsequently, the sample is exposed to means for differentially counting hybridization beads in order to quantify first and second hybridization probes which have formed duplexes. The numbers obtained provide a precise estimate of the number of copies of the reference and target polynucleotide in the population because differential counting

30 means count individual beads. One bead is equal to one probe which, in turn, signifies one copy of the nucleic acid being measured.

- 20 -

An example of a differential counting means is an impedance measuring device, such as a Coulter counter (Coulter Electronics, Inc., Miami, Florida). Sample is passed through the device which differentially detects the two types of hybridization beads by measuring their differential impedance of an electric current. Alternatively, the device
5 may measure fluorescence, color, or other parameters. In order to increase the speed of the assay, a multi-orifice device may be used. A multi-orifice impedance counter is shown schematically in figure 2. A multi-orifice array is placed at one end of a column filled with an electrically-conductive fluid, such as saline. Hybridization beads with either hybridized target or reference segments are inserted at an opposite end of the
10 column. Each orifice is large enough to accommodate only one hybridization bead at a time and sufficiently wide to allow reliable impedance measurements. A voltage is set across each orifice. Each hybridization bead (which is non-conducting), as it passes through one of the orifices, displaces a volume of saline, thus creating an impedance that is proportional to its size. This, in turn, creates a measurable decrease in current
15 that is directly correlated with the size of the bead. By compiling the number of each of the two distinct impedance events, a precise estimate of the number of hybridization beads and, therefore, the number of probes of each type in the population may be obtained.

Upon quantitative measurement of first and second hybridization beads, the data
20 may be analyzed to determine whether any difference between the amounts of first and second hybridization beads is statistically significant. A reduction in the amount of target relative to the reference is indicative of a mutation in or deletion of the target allele in a subpopulation of cells in the sample. Where the p53 gene is the target allele, such a mutation is indicative of a cancerous or precancerous condition. A
25 clinician may use such results as a basis for prescribing additional treatment, such as endoscopy and polypectomy procedures.

B. Detection of Mutations in Single-base Polymorphisms

The basic method described above may also be applied to detect a loss of heterozygosity or other mutation at a single base polymorphic site between maternal
30 and paternal alleles. Such detection is typically an indication of a larger deletion or other mutation. However, a mutation at a single polymorphic nucleotide may be all that

- 21 -

is necessary to inhibit gene function in one of the two alleles. A mutation in a single-base polymorphic region may be difficult to detect due to a recently-discovered phenomenon called complementary reduplication. In complementary reduplication, the loss of one of two alleles at a particular locus results in "reduplication" of the surviving allele. Reduplication usually takes place on the chromosome containing the surviving allele and involves the production of one or more copies of the surviving allele in close proximity on the chromosome to the position of the surviving allele. In the case of a locus that displays one or more single-base allelic polymorphisms (*i.e.*, heterozygosity at the locus is determined by virtue of one or more single-base differences in one or more regions of the locus), complementary reduplication results in the insertion on the chromosome containing the surviving allele of a duplicate of the sequence corresponding to that which was deleted. Even under the most stringent hybridization conditions, some of a probe directed against the deleted sequence will bind to the reduplicated sequence at a locus of a single-base polymorphism. Accordingly, in such circumstances, the deletion may not be detected because any true difference in the number of probes binding to the polymorphic site (*i.e.*, the allelic region encompassing the single-base polymorphism) may be obscured by an increase resulting from the other allele's reduplicated region.

The problems associated with complementary reduplication, and with non-specific probe binding generally, are alleviated by the practice of the methods described herein. Such methods allow detection of a deletion in one of two alleles present at a specific locus in a subpopulation of cells contained in a biological sample. Numerous alleles, including tumor suppressor alleles, contain single polymorphic nucleotides in the context of a constant nucleic acid region. Individuals normally may be either homozygous or heterozygous for the polymorphic nucleotide. Since numerous single-base polymorphic nucleotide sites exist in most alleles, the probability that a given individual is heterozygous at least one of the single-base polymorphism sites is high. A statistically-significant reduction in one of the two nucleotides at a single-base polymorphic site (at which the individual is heterozygous) may be used as a marker for a deletion in the allele encompassing that site.

- 22 -

Genomic regions containing known single-base polymorphisms may be identified by reference to a nucleotide database, such as GenBank, EMBL, or any other appropriate database. The existence of polymorphisms may be determined by methods taught herein, gel electrophoresis or by other standard methods. For
5 purposes of the invention, a single-base polymorphism is intended to be a single polymorphic nucleotide adjacent to a non-polymorphic region of the allele regardless of whether the single polymorphic nucleotide forms part of a larger polymorphic site (i.e. the single-base polymorphism may be the terminal nucleotide of a larger,
polynucleotide polymorphism). For cancer detection, the regions considered are
10 regions in which loss of heterozygosity is prevalent, such as regions containing tumor suppressor genes. A given individual may be homozygous or heterozygous for the polymorphic nucleotide in any identified single-base polymorphic region. Accordingly, if a number of single-base polymorphic regions are identified, the probability increases that at least one heterozygous single-base polymorphic region is found in a sample.

15 Once single-base polymorphic sites are identified, a DNA sample is obtained from a patient, e.g., from blood cells, to determine which of those sites is heterozygous in normal (i.e., non-cancerous or non pre-cancerous) cells for that individual. Then, a stool sample is prepared as described above. Double stranded DNA in the sample is converted to single-stranded DNA. Then, either the coding strand or the anti-coding
20 strand for both alleles is removed from the sample. As will be evident from the following discussion, methods disclosed herein are indifferent as to whether coding strand or anti-coding strand is tested.

An oligonucleotide probe is constructed that is complementary to a portion of the region of single-base polymorphism, said portion ending at the nucleotide that is
25 immediately 3' to the polymorphic nucleotide, regardless of whether the 5'-3' (coding) strand or the 3'-5' (anticoding) strand is used as a template. Figure 3 shows four possible probes that are immediately 3' to the polymorphic nucleotide for each of four possible template strands as described above (the Sequences in Figure 3 are hypothetical and are not intended to represent any actual sequence). The sequence
30 labeled M1 is SEQ ID NO:1; the sequence labeled M2 is SEQ ID NO:2; the sequence

- 23 -

labeled M3 is SEQ ID NO: 3; the sequence labeled M4 is SEQ ID NO:4; the sequence labeled F1 is SEQ ID NO: 5; the sequence labeled F2 is SEQ ID NO: 6; the sequence labeled F3 is SEQ ID NO: 7; and the sequence labeled F4 is SEQ ID NO: 8. While either strand may be used as a template for probe binding to determine heterozygosity and/or the loss thereof, the sequence of the probe that is hybridized to the template will be different depending upon the strand used. Probes may be of any length that allows efficient and specific hybridization. Figure 3 merely illustrates four hypothetical probes that are useful for hybridization to the hypothetical sequence shown. The length of probe sequences may be determined as appropriate for each genomic region that is analyzed. A preferable length is between about 10 and about 100 nucleotides. The size of the probe will also depend upon the size of the region surrounding the single-base polymorphism (*i.e.*, the region 5' or 3' to the next adjacent polymorphism, if any). Details concerning the construction and hybridization of oligonucleotide probes are known in the art.

Unique probes for each polymorphic region will hybridize to regions of both maternal and paternal alleles up to, but not including, the polymorphic nucleotide, which, in a heterozygote, will be different in the maternal and paternal alleles. Figure 3 shows only a small portion of the region surrounding the polymorphic nucleotide. The alleles shown in Figure 3 are heterozygous at the polymorphic site.

Probe is hybridized to its specific template DNA by standard methods. The sample may optionally be washed to remove unhybridized probe. To determine whether each target region bound by a probe is heterozygous or homozygous at the polymorphic nucleotide, a modification of the dideoxy chain termination method as reported in Sanger, *Proc. Nat'l Acad. Sci. (USA)*, 74: 5463-5467 (1977), incorporated by reference herein, is used. The method involves using at least two of the four common 2', 3'-dideoxy nucleoside triphosphates (ddATP, ddCTP, ddGTP, and ddTTP). A different detectable label is attached to each dideoxy nucleoside triphosphate (ddNTP) according to methods known in the art. Differentially-labeled ddNTPs are available commercially, for example, from Perkin Elmer Corporation (Cat. No. 401456). At least two labeled ddNTPs then are exposed to each sample having probe hybridized to maternal and paternal alleles as described above. The choice of which two ddNTPs

- 24 -

are used will depend upon the nucleotides at the heterozygous polymorphic site. Any 3' modified nucleoside triphosphate may be used in the method as long as the 3' modification prevents binding of an additional 3' nucleotide (i.e. probe extension) and does not inhibit binding of the modified nucleotide to the 3' end of the probe. A DNA
5 polymerase, such as Sequenase™ (Perkin-Elmer), is added to the sample mixture. Using the allelic strands as primer, the polymerase will add one ddNTP to the 3' end of the probe, the incorporated ddNTP will be complementary to the nucleotide that exists at the single-base polymorphic site. Because the ddNTPs have no 3' hydroxyl, further elongation of the hybridized probe will not occur. After completion, the sample is
10 washed to remove excess ddNTPs. Label is then counted in each sample. The presence of two differentially-labeled ddNTPs in a sample is indicative of heterozygosity at the polymorphic site.

It is not necessary to determine the amount of each label present in the sample in order to establish heterozygosity or homozygosity. For Example, differentially-
15 labeled deoxynucleoside triphosphates may be used for a determination of heterozygosity or homozygosity. The mere fact that two different labeled dideoxy nucleotides are incorporated into the probe means that the single-base polymorphic site being analyzed is heterozygous. However, determination of sites at which a patient is polymorphic is useful in order to establish a baseline of polymorphisms which
20 may be used in future tests to detect changes in polymorphic sites which may be indicative of cancer. The existence of polymorphisms may be determined by methods taught herein, by gel electrophoresis or by other standard methods.

In the case in which heterozygosity exists at the polymorphic site, counting the amount of each of the two differentially-labeled ddNTPs allows a determination of
25 whether there is a loss of heterozygosity (i.e., a deletion) in a subpopulation of cells in the sample. In a normal (i.e., non-cancerous) sample containing cells that are heterozygous at the single-base polymorphic site, it is expected that the detected amount of each of the two ddNTPs added to the probe will be identical (within chosen limits of statistical significance). However, if a deletion has occurred in one of the two
30 alleles in a subpopulation of cells in the sample, there will be a statistically-significant difference between the amounts of each of the two alleles detected via the incorporated

- 25 -

(labeled) ddNTPs. The detection of such a difference is indicative of genomic instability within the sample. Such genomic instability indicates the possibility of cancerous or pre-cancerous cells in the sample.

5 In order to improve the ability to count alleles to which ddNTPs have attached accurately, ddNTPs are labeled with hybridization-type beads of different sizes as described above. Alleles with bound probe comprising a labeled ddNTP are counted as described above using a counting device, such as a Coulter counter. Also as described above, differential fluorescent labels or other counting means may be used to separately detect incorporated ddNTPs.

10 The detection of heterozygosity at single-base polymorphic sites and the detection of the loss of heterozygosity may be determined in separate steps. For example, probes may be hybridized immediately adjacent to but not including the nucleotide determined to be polymorphic as described above. The four ddNTPs may then be added to the sample, washed, and the presence or absence of each label may
15 be detected. Detection of only one label indicates that the individual from whom the sample was obtained is homozygous at the site of the potential polymorphic nucleotide. Detection of two labels means that the individual is heterozygous. The heterozygous loci are recorded. As noted above, baseline determinations of heterozygosity may be done using standard methods. Once a baseline is established, future tests on that
20 individual are performed exploiting the heterozygous loci in order to detect a loss of heterozygosity. For the detection of cancer, the heterozygous loci are typically chromosomal areas containing tumor suppressor genes, including p53, dcc, apc, and others. Using methods described herein, a "fingerprint" of heterozygous tumor suppressor loci may be constructed. Future deviation from the fingerprint (*i.e.*,
25 deletions) provides valuable information as to the development of cancer.

A preferred use of the foregoing methods is in the detection of colon cancer. A representative stool sample is prepared as described above. Double-stranded DNA is converted to single-stranded DNA and complement of the strand to be detected is removed from the sample. The remaining single-stranded DNA is exposed to multiple
30 copies of a probe designed on the basis of known single-base polymorphisms in a cancer-associated allele such that the probe hybridizes with a desired number of

- 26 -

nucleotides immediately adjacent the polymorphic nucleotide as described above. After hybridization is complete, the sample is washed and exposed to differentially-labeled ddNTPs and a DNA polymerase. The sample then is washed to remove unincorporated ddNTPs. The presence of any labeled ddNTPs is determined. If two
5 labels are detected, the individual from whom the sample is obtained is heterozygous at the polymorphic nucleotide. The heterozygosity of the allele and the probe sequence matching the site immediately adjacent to the polymorphic allele are noted for reference in future testing for the loss of heterozygosity. Alternatively, once the patient is determined to be heterozygous at a locus, an assay may be performed
10 immediately in the manner described above in order to determine an existing loss of heterozygosity in a subpopulation of cells in the sample.

C. Analysis of Microsatellite Instability

Microsatellites are di- or trinucleotide repeats found throughout the genome. A particular array of microsatellite repeats is often associated with a particular genomic
15 sequence and is stably inherited under normal conditions. Expansions of microsatellite copy number typically, called "microsatellite instability," are associated with defects in mismatch repair. Accordingly, changes in a microsatellite region indicate that the patient is at risk for a mutation in other genomic regions.

In order to detect microsatellite instability as an indicator of a mutation in a
20 cancer-associated gene, one must first identify a microsatellite region associated with the gene of interest. Such regions are typically identified on a database, such as GenBank, EMBL, and others. Once a wild-type microsatellite region associated with, for example, the p53 tumor suppressor gene, is identified, an oligonucleotide probe is constructed that spans the microsatellite region and the regions immediately 5' and
25 immediately 3' to the microsatellite region. The precise length of probes may be determined by the experimenter. Probes are constructed that hybridize to the microsatellite region, including portions extending 5' and 3', on both the maternal and paternal alleles with which the microsatellite is associated (e.g., p53).

An appropriate sample of body tissue or fluid is obtained and processed as
30 described herein. Double stranded DNA is denatured and an excess of maternal and

- 27 -

paternal probes, as described above, are introduced into the sample under hybridization conditions. The probes are detectably labeled as described above. Complement of the strands to be detected may optionally be removed by methods described above. The sample is then washed to remove unhybridized probe and the amount of hybridized probe is quantitatively detected.

Quantitative detection may be accomplished by any means described herein. For example, probes may be attached to hybridization beads such that probes that bind to maternal allele are attached to beads of one size and probes that bind to paternal allele are attached to beads of a second size that is distinguishable from beads of the first size. Beads with attached probe may be counted as described above.

The detection of a statistically-significant difference between the amount of probe binding to the maternal allele and the amount of probe binding to the paternal allele is indicative of microsatellite instability. As previously mentioned, microsatellite instability can be indicative of a mutation at the locus in which the microsatellite resides. If the microsatellite region is associated with a tumor suppressor gene or an oncogene, the detection of microsatellite instability in an allele in a subpopulation of cells in a biological sample is indicative of the potential for cancer or that cancer or precancer may have already developed. Further testing as described herein (either by invasive or noninvasive means) may then be conducted.

In an alternative embodiment, a "fingerprint" of microsatellites is taken from regions associated with cancer-causing genes in a sample obtained from a patient. Such a fingerprint may be obtained by standard methods. The fingerprint comprises the sequence of wild-type microsatellites associated with the cancer-causing gene or genes. Once obtained, the fingerprint is stored and is used in future tests of samples from the same patient in order to monitor changes in microsatellite regions (i.e. microsatellite instability) that may be associated with the development of cancer. Changes in microsatellite length and/or sequence over time may be used to prescribe additional testing and/or treatment in order to detect and remove cancerous tissue at an early stage in its etiology.

Additional embodiments of the invention are apparent upon consideration of the following claims.

- 28 -

CLAIMS

What is claimed is:

- 1 1. A method of screening for the presence of a disease in a patient, the method
2 comprising the steps of:
 - 3 a) obtaining a sample comprising a circumferential surface of a stool
4 voided by a patient; and
 - 5 b) performing an assay to detect in the sample characteristics
6 indicative of the presence of said disease.
- 1 2. A method for screening for the presence of a colorectal cancer or precancerous
2 lesion in a patient, the method comprising the steps of:
 - 3 a) obtaining a stool voided by a patient;
 - 4 b) removing a circumferential surface from said stool; and
 - 5 c) performing an assay on said surface to detect characteristics
6 indicative of the presence of cells or cellular debris shed onto said
7 stool.
- 1 3. The method of claim 2, wherein said assay detects debris from a clonal
2 population of transformed cells comprising said lesion.
- 1 4. The method of claim 3, wherein said assay detects a protein expressed by the
2 transformed cells indicative of the presence of said cells.
- 1 5. The method of claim 3, wherein said assay detects a DNA characteristic
2 indicative of the presence of said cells.
- 1 6. The method of claim 1, further comprising the step of homogenizing said portion
2 in a physiologically compatible buffer prior to step (b).
- 1 7. The method of claim 6, wherein said physiologically compatible buffer comprises
2 a detergent and a proteinase.
- 1 8. The method of claim 1, wherein said assay detects the presence of
2 carcinoembryonic antigen shed from said lesion.

- 29 -

- 1 9. The method of claim 1, wherein said assay comprises the step of exposing said
2 sample to an antibody which specifically binds a molecule characteristic of the
3 presence of said debris.
- 1 10. The method of claim 5, wherein said DNA characteristic is a mutation.
- 1 11. The method of claim 10, wherein said mutation is selected from the group
2 consisting of loss of heterozygosity and microsatellite instability.
- 1 12. The method of claim 10, wherein said characteristics comprise a deletion in a
2 tumor suppressor allele.
- 1 13. The method of claim 1, wherein said assay comprises the step of determining
2 whether a difference exists in said sample between a number X of a first allele
3 known or suspected to be mutated in a subpopulation of cells in the sample and
4 a number Y of a second allele that is known or suspected not to be mutated in a
5 subpopulation of cells in the sample, the presence of a statistically-significant
6 difference being indicative of a mutation in a subpopulation of cells in the
7 sample and the potential presence of a cancerous or precancerous lesion.
- 1 14. The method of claim 1, wherein said assay comprises the step of determining
2 whether a difference exists between a number of a target tumor suppressor
3 allele in the sample and a number of a non-cancer-associated reference allele in
4 the sample, the presence of a statistically-significant difference being indicative
5 of a deletion of the target tumor suppressor allele in a subpopulation of cells in
6 the sample and the potential presence of a cancerous or precancerous lesion.
- 1 15. The method of claim 5, wherein said characteristics comprise a loss of
2 heterozygosity encompassing a polymorphic locus.
- 1 16. The method of claim 13, wherein said assay comprises the steps of:
2 a) detecting an amount of a maternal allele at a polymorphic locus in
3 the biological sample;
4 b) detecting an amount of a paternal allele at the polymorphic locus in
5 the biological sample; and

- 30 -

- 6 c) determining whether a difference exists between the amounts of
7 maternal and paternal allele,
8 the presence of a statistically-significant difference being indicative of a deletion
9 at the polymorphic locus in a subpopulation of cells in the biological sample and
10 the potential presence of a lesion.
- 1 17. The method of claim 1, comprising the additional step of performing a visual
2 examination of the colon of a patient presenting positive assay results.
- 1 18. The method of claim 1, wherein said at least cross-sectional portion comprises
2 an entire stool voided by a patient.
- 1 19. A method for reducing morbidity from colorectal cancer in a population, the
2 method comprising the steps of:
- 3 a) obtaining a sample comprising at least a representative portion of
4 a stool voided by a patient;
5 b) performing an assay to detect in the sample the presence of
6 cellular debris shed from the lesion into the voided stool; and
7 c) performing a visual examination of the colon of a patient presenting
8 positive results in said assay to detect the presence of a lesion.
- 1 20. The method of claim 19, wherein said assay detects a loss of DNA in a portion of
2 debris shed from a clonal population of transformed epithelial cells comprising
3 said lesion.




FIG. 1




FIG. 3

4/4

1: 3'ACGCTACGG5'
2: 5'....ATCGGCTTA**C**TGCGATGCC....3'

M

3: 3'....TAGCCGAAT**G**ACGCTACGG....5'
4: 5'ATCGGCTTA3'

1: 3'ACGCTACGG5'
2: 5'....ATCGGCTTA**T**TGCGATGCC....3'

F

3: 3'....TAGCCGAATA**A**ACGCTACGG....5'
4: 5'ATCGGCTTA3'

FIG. 4**SUBSTITUTE SHEET (RULE26)**

SEQUENCE LISTING

<110> Lapidus, Stanley N
Shuber, Anthony P
EXACT LABORATORIES

<120> Methods for Disease Diagnosis From Stool Samples

<130> EXT-015PC

<140>
<141>

<150> US 09/059,713
<151> 1998-04-13

<160> 8

<170> PatentIn Ver. 2.0

<210> 1
<211> 9
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
M1

<400> 1
ggcatcgca 9

<210> 2
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
M2

<400> 2
atcggcttac tgcgatgcc 19

<210> 3
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
M3

<400> 3
ggcatcgag taagccgat 19

<210> 4
<211> 9
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
M4

<400> 4
atcggctta 9

<210> 5
<211> 9
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
F1

<400> 5
ggcatcgca

9

<210> 6
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
F2

<400> 6
atcggcttat tgcgatgcc

19

<210> 7
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
F3

<400> 7
ggcatcgcaa taagccgat

19

<210> 8
<211> 9
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Hypothetical
F4

<400> 8
atcggctta

9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/08013

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9728450	A	07-08-1997	US 5741650 A	21-04-1998
			AU 704696 B	29-04-1999
			AU 2240697 A	22-08-1997
			CA 2215263 A	07-08-1997
			EP 0817968 A	14-01-1998
EP 391565	A	10-10-1990	US 5192501 A	09-03-1993
			CA 2012830 A	04-10-1990
			DE 69026859 D	13-06-1996
			DE 69026859 T	12-09-1996
			JP 2295495 A	06-12-1990
			US 5196167 A	23-03-1993
			US 5217874 A	08-06-1993
US 4578358	A	25-03-1986	AU 547546 B	24-10-1985
			AU 2470984 A	08-11-1984
			BR 8400933 A	05-03-1985
			CA 1212891 A	21-10-1986
			EP 0124215 A	07-11-1984
			JP 1688475 C	11-08-1992
			JP 3054305 B	19-08-1991
			JP 59204766 A	20-11-1984
			ZA 8401079 A	26-09-1984